

1 581 Rec'd PC 19 DEC 2001

DESCRIPTION

PHTHALAMIDE DERIVATIVES, INTERMEDIATES IN THE
PRODUCTION THEREOF, AND AGRICULTURAL/HORTICULTURAL
INSECTICIDES AND METHOD FOR USING THE SAME

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

The present invention relates to phthalamide derivatives, production intermediates thereof, agrohorticultural insecticides containing said
5 compounds as active ingredient, and a method for using said insecticides.

RELATED ART

A part of the phthalamide derivatives of the present invention are disclosed in JP-A-59-163353, JP-
10 A-61-180753, Journal of Chemical Society (J.C.S.), Perkin I, 1338-1350 (1978), etc. Neither description nor suggestion about usefulness of these compounds as an agrihorticultural insecticide, however, is made therein at all. On the other hand, the heterocyclic
15 amine derivatives represented by the general formula (IV), which serve as intermediate compounds for production of said phthalamide derivatives, are novel compounds not found in literature.


SUMMARY OF THE INVENTION

20 The present inventors have conducted repeated

studies on the development of a novel agrohorticultural agent. As a result, it has been found that the phthalamide derivatives of the present invention represented by the general formula (I) which are novel compounds and some known compounds disclosed in prior art are useful as novel agrohorticultural insecticides. It has further been found that the heterocyclic amine derivatives represented by the formulas (IV') which are novel compounds not found in literature are useful as intermediates for production of a variety of physiologically active compounds usable as medical drugs, pesticides, etc. Based on these findings, the present invention has been accomplished.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to phthalamide derivatives represented by the general formula (I):



{wherein R^1 , R^2 and R^3 , which may be same or different, represent hydrogen atom, C_3 - C_6 cycloalkyl group, halo C_3 - C_6 cycloalkyl group or $-A^1-(G)_r$ (in this formula, A^1 represents C_1 - C_8 alkylene group, C_3 - C_6 alkenylene group or C_3 - C_6 alkynylene group; G , which may be same or different, represents hydrogen atom, halogen atom,

cyano group, nitro group, halo C₁-C₆ alkyl group, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, C₁-C₆ alkoxy carbonyl group, di(C₁-C₆) alkoxyphosphoryl group in which the (C₁-C₆) alkoxy groups may be same or different, di(C₁-C₆) alkoxythiophosphoryl group in which the (C₁-C₆) alkoxy groups may be same or different, diphenylphosphino group, diphenylphosphono group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (as used herein, the term "heterocyclic group" means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl group), substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkyl-

sulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆
 alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group,
 or -Z³-R⁴ (in this formula, Z³ represents -O-, -S-, -SO-,
 -SO₂-, -N(R⁵)- (in this formula, R⁵ represents hydrogen
 5 atom, C₁-C₆ alkylcarbonyl group, halo C₁-C₆ alkylcarbonyl
 group, C₁-C₆ alkoxy carbonyl group, phenylcarbonyl group,
 substituted phenylcarbonyl group having at least one,
 same or different substituents selected from the group
 consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆
 10 alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group,
 C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆
 alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group,
 C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl
 group, phenyl C₁-C₄ alkoxy carbonyl group, substituted
 15 phenyl C₁-C₄ alkoxy carbonyl group having, on the ring
 thereof, at least one, same or different substituents
 selected from the group consisting of halogen atom,
 C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy
 group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group,
 20 halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group,
 halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group
 and halo C₁-C₆ alkylsulfonyl group, C₁-C₆ alkylsulfonyl
 group or halo C₁-C₆ alkylsulfonyl group), -C(=O)- or
 -C(=NOR⁶)- (in this formula, R⁶ represents hydrogen
 25 atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₃-C₆
 alkenyl group, halo C₃-C₆ alkenyl group, C₃-C₆ alkynyl
 group, C₃-C₆ cycloalkyl group, phenyl C₁-C₄ alkyl group,
 or substituted phenyl C₁-C₄ alkyl group having, on the

ring thereof, at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group), and R⁴ represents hydrogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₃-C₆ alkenyl group, halo C₃-C₆ alkenyl group, C₃-C₆ alkynyl group, halo C₃-C₆ alkynyl group, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, C₁-C₆ alkoxy C₁-C₆ alkyl group, C₁-C₆ alkylthio C₁-C₆ alkyl group, formyl group, C₁-C₆ alkylcarbonyl group, halo C₁-C₆ alkylcarbonyl group, C₁-C₆ alkoxycarbonyl group, mono (C₁-C₆) alkylaminocarbonyl group, di(C₁-C₆) alkylaminocarbonyl group in which the (C₁-C₆) alkyl groups may be same or different, mono(C₁-C₆) alkylaminothiocarbonyl group, di(C₁-C₆) alkylaminothiocarbonyl group in which the (C₁-C₆) alkyl groups may be same or different, di(C₁-C₆) alkoxyphosphoryl group in which the (C₁-C₆) alkoxy groups may be same or different, di(C₁-C₆) alkoxythiophosphoryl group in which the (C₁-C₆) alkoxy groups may be same or different, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkyl-

- sulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, phenyl C₁-C₄ alkyl group, substituted phenyl (C₁-C₄) alkyl group having, on the ring thereof, at least one,
- 5 same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group,
- 10 C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), or substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents
- 15 selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group); and r represents an
- 20 integer of 1 to 4); further, R¹ and R² may be taken conjointly to form 4- to 7-membered rings which may be intercepted by 1 to 3, same or different oxygen atom, sulfur atom or nitrogen atom;
- 25 X, which may be same or different, represents halogen atom, cyano group, nitro group, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, phenyl group, substituted phenyl group having at least one, same or

different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, or -A²-R⁷ [in this formula, A² represents -O-, -S-, -SO-, -SO₂-, -NR⁸- (in this formula R⁸ represents hydrogen atom, C₁-C₆ alkylcarbonyl group, halo C₁-C₆ alkylcarbonyl group, C₁-C₆ alkoxy carbonyl group, phenylcarbonyl group, substituted phenylcarbonyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, phenyl C₁-C₄ alkoxy carbonyl group or substituted phenyl

- C₁-C₄ alkoxy carbonyl group having, on the ring thereof, at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group), -C(=O)-, -C(=NOR⁶)- (in this formula, R⁶ is as defined above), C₁-C₆ alkylene group, halo C₁-C₆ alkylene group, C₂-C₆ alkenylene group, halo C₂-C₆ alkenylene group, C₂-C₆ alkynylene group or halo C₃-C₆ alkynylene group; and
- (1) in cases where A² represents -O-, -S-, -SO-, -SO₂- or -NR⁸- (in this formula, R⁸ is as defined above), R⁷ represents hydrogen atom, halo C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkenyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group,

halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, or -A³-R⁹ (in this formula, A³ represents C₁-C₆ alkylene group, halo C₁-C₆ alkylene group, C₃-C₆ alkenylene group, halo C₃-C₆ alkenylene group, C₃-C₆ alkynylene group or halo C₃-C₆ alkynylene group; and R⁹ represents hydrogen atom, halogen atom, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, C₁-C₆ alkoxycarbonyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, or -A⁴-R¹⁰ (in this formula, A⁴ represents -O-, -S-, -SO-, -SO₂- or -C(=O)-; and R¹⁰ represents C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₃-C₆ alkenyl group, halo C₃-C₆ alkenyl group, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl

group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), or substituted heterocyclic group (the
 5 term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio
 10 group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group));

(2) in cases where A² represents -C(=O)- or -C(=NOR⁶)- (in this formula, R⁶ is as defined above), R⁷
 15 represents hydrogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₂-C₆ alkenyl group, halo C₂-C₆ alkenyl group, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, mono(C₁-C₆) alkylamino group, di(C₁-C₆) alkylamino group
 20 in which the (C₁-C₆) alkyl groups may be same or different, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy
 25 group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, phenylamino group,

substituted phenylamino group having, on the ring thereof, at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), or substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group; and

(3) in cases where A² represents C₁-C₆ alkylene group, halo C₁-C₆ alkylene group, C₂-C₆ alkenylene group, halo C₂-C₆ alkenylene group, C₂-C₆ alkynylene group or halo C₃-C₆ alkynylene group, R⁷ represents hydrogen atom, halogen atom, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, C₁-C₆ alkoxycarbonyl group, tri(C₁-C₆) alkylsilyl group in which the (C₁-C₆) alkyl groups may be same or different, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen

atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, or -A⁵-R¹¹ (in this formula, A⁵ represents -O-, -S-, -SO- or -SO₂-; and R¹¹ represents C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different

substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, or -A⁶-R¹² (in this formula, A⁶ represents C₁-C₆ alkylene group, halo C₁-C₆ alkylene group, C₂-C₆ alkenylene group, halo C₂-C₆ alkenylene group, C₂-C₆ alkynylene group or halo C₃-C₆ alkynylene group; and R¹² represents hydrogen atom, halogen atom, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group, halo C₁-C₆ alkylsulfonyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, phenoxy group, substituted phenoxy group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkyl-







sulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, phenylthio group, substituted phenylthio group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), or substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group))];


n represents an integer of 0 to 4; further, X may be taken conjointly with the adjacent carbon atom on the phenyl ring to form a fused ring (as used herein, the term fused ring means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene,

benzoxazole, benzothiazole, benzimidazole or indazole), and said fused ring may have at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group, halo C₁-C₆ alkylsulfonyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), and substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group;


Q represents an N-, S- or O-containing, optionally substituted, heterocyclic group or fused heterocyclic group, selected from the group consisting

of the following formulas Q1 to Q60;







Q34




Q35




Q36




Q37




Q38




Q39




Q40




Q41




Q42




Q43




Q44




Q45




Q46




Q47




Q48




Q49




Q50




Q51




Q52




Q53




Q54




Q55




Q56




Q57



Q58



Q59



Q60

(in these formulas, Y, which may be same or different, represents halogen atom, cyano group, nitro group, halo C₃-C₆ cycloalkyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group is as defined above), substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, or -A²-R⁷ (in this formula, A² and R⁷ are as defined above); m represents an integer of 0 to 6; R¹³ in the formula Q22 and Q23 represents hydrogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₃-C₆ alkenyl group, halo C₃-C₆ alkenyl group, C₃-C₆ alkynyl group, halo C₃-C₆ alkynyl group, C₃-C₆ cycloalkyl group, halo C₃-C₆ cycloalkyl group, C₁-C₆ alkoxy C₁-C₆ alkyl group, halo C₁-C₆ alkoxy C₁-C₆ alkyl group, C₁-C₆ alkylthio C₁-C₆ alkyl group, halo C₁-C₆ alkylthio C₁-C₆

- alkyl group, C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group, halo C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group, C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group, halo C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group, C₁-C₆ alkylsulfonyl group, halo C₁-C₆ alkylsulfonyl group, C₁-C₆ alkylcarbonyl group, halo C₁-C₆ alkylcarbonyl group, C₁-C₆ alkoxy carbonyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, phenyl C₁-C₄ alkyl group, substituted phenyl C₁-C₄ alkyl group having, on the ring thereof, at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, phenylcarbonyl group, or substituted phenylcarbonyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆

alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group);

alternatively, Y may be taken conjointly with adjacent carbon atom on the ring to form a fused ring (the fused ring is as defined above), and said fused
5 ring may have at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl
10 group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group, halo C₁-C₆ alkylsulfonyl group, phenyl group, substituted phenyl group having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆
15 alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group, heterocyclic group (the term heterocyclic group
20 is as defined above), and substituted heterocyclic group (the term heterocyclic group is as defined above) having at least one, same or different substituents selected from the group consisting of halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy
25 group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group and halo C₁-C₆ alkylsulfonyl group;

W represents O, S or N-R¹³ (in this formula, R¹³ is as defined above); and Z¹ and Z² represent oxygen atom or sulfur atom;

provided that when X, R¹ and R³ simultaneously represent hydrogen atom, Z¹ and Z² simultaneously represent oxygen atom, Q represents Q27, and Y is a chlorine atom of 2-position, then R² is not 1,2,2-trimethylpropyl group};


and to an agrohorticultural insecticide, and a method for using the same.

The present invention further relates to a heterocyclic amine derivative represented by the following general formula (IV'):



wherein:



(1) in cases where Q' represents one of the following formulas Q26, Q28-Q31 and Q33-Q39,



Y, which may be same or different, represents hydrogen atom, halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group or halo C₁-C₆ alkylsulfonyl group, m represents an integer of 1 to 4, and at least one of Y, of which total number is m, is perfluoro C₂-C₆ alkyl group;

10 and

(2) in a case where Q' represents Q27 and Q32:



Y, which may be same or different, represents hydrogen atom, halogen atom, C₁-C₆ alkyl group, halo C₁-C₆ alkyl group, C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy group, C₁-C₆ alkylthio group, halo C₁-C₆ alkylthio group, C₁-C₆ alkylsulfinyl group, halo C₁-C₆ alkylsulfinyl group, C₁-C₆ alkylsulfonyl group or halo C₁-C₆ alkylsulfonyl group, m represents an integer of 1 to 4, and at least one of Y, of which total number is m, is perfluoro C₂-C₆ alkyl group, halo C₁-C₆ alkoxy group, halo C₁-C₆ alkoxy halo C₁-C₆ alkoxy group or halo C₁-C₆ alkylthio group.

The heterocyclic amine derivative of the formula (IV') is useful for an intermediate compound for production of the phthalamide derivatives of the

formula (I).

In the definition of the general formula (I) of the phthalamide derivative of the present invention, "halogen atom" means chlorine atom, bromine atom, iodine atom or fluorine atom; "C₁-C₆ alkyl" means a straight or branched chain alkyl group having 1 to 6 carbon atoms such as methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, n-hexyl or the like; "halo C₁-C₆ alkyl" means a straight or branched chain alkyl group having 1 to 6 carbon atoms substituted with at least one, same or different halogen atoms; and "C₁-C₈ alkylene" means a straight or branched chain alkylene group having 1 to 8 carbon atoms such as methylene, ethylene, propylene, trimethylene, dimethylmethylene, tetramethylene, isobutylene, dimethylethylene, octamethylene or the like.

As examples of the "R¹ and R² taken conjointly to form a 4- to 7-membered ring which may be intercepted by 1 to 3, same or different oxygen atom, sulfur atom or nitrogen atom", there can be referred to azetidine ring, pyrrolidine ring, pyrroline ring, piperidine ring, imidazolidine ring, imidazoline ring, oxazolidine ring, thiazolidine ring, isoxazolidine ring, isothiazolidine ring, tetrahydropyridine ring, piperazine ring, morpholine ring, thiomorpholine ring, dioxazine ring, dithiazine ring, and the like.

Some of the phthalamide derivatives of the


present invention represented by the general formula (I) have an asymmetric carbon atom or asymmetric center in the structural formulas thereof, and there can exist two optical isomers sometimes. The present invention
 5 includes all such optical isomers and their mixtures at arbitrary proportions, and sometimes includes salts and hydrates thereof.

In the phthalamide derivatives of the present invention represented by the general formula (I),
 10 preferable substituents are as follows. Thus, the phthalamide derivative of the invention is preferably a phthalamide derivative of the general formula (I) wherein R^1 , R^2 and R^3 may be same or different and represent hydrogen atom or $-A^1-G$ (in this formula, A^1
 15 represents C_1-C_8 alkylene group and G represents hydrogen atom, C_1-C_6 alkylthio group, C_1-C_6 alkylsulfinyl group, C_1-C_6 alkylsulfonyl group, C_1-C_6 alkylcarbonyl-amino group or C_1-C_6 alkoxycarbonylamino group); X may be same or different and represents halogen atom, nitro
 20 group, halo C_1-C_6 alkyl group, halo C_1-C_6 alkoxy group or halo C_1-C_6 alkylthio group; n represents an integer of 0 to 4; Q represents Q27; Y may be same or different and represents halogen atom, C_1-C_6 alkyl group, halo C_1-C_6 alkyl group, C_1-C_6 alkoxy group, halo C_1-C_6 alkoxy group,
 25 halo C_1-C_6 alkoxy halo C_1-C_6 alkoxy group, C_1-C_6 alkylthio group, halo C_1-C_6 alkylthio group, halo C_1-C_6 alkoxy halo C_1-C_6 alkylthio group, halo C_1-C_6 alkylsulfinyl group or halo C_1-C_6 alkylsulfonyl group; m represents an integer

of 0 to 4; Z^1 and Z^2 represent oxygen atom. Further preferably, the phthalamide derivative of the invention is a phthalamide derivative of general formula (I) wherein R^1 and R^3 represent hydrogen atom; R^2 represents C_1-C_6 alkylthio C_1-C_6 alkyl group; X represents halogen atom; n represents an integer of 1 to 2; Q represents Q27; Y may be same or different and represents halogen atom, C_1-C_6 alkyl group, halo C_1-C_6 alkyl group, C_1-C_6 alkoxy group or halo C_1-C_6 alkoxy group; m represents an integer of 1 to 2; and Z^1 and Z^2 represent oxygen atom.

The compounds of the present invention can be produced according to Schemes 1 and 2 mentioned below, though the compounds of the present invention can also be produced by the process described in JP-A-11-240857.

15 Production process 1



wherein R^1 , R^2 , Z^1 , Z^2 , X, Q and n are as defined above.

A phthalic anhydride derivative represented by general formula (V) is reacted with a heterocyclic amine derivative represented by general formula (IV) in the presence of an inert solvent to form a phthalimide derivative represented by general formula (III). After isolating or without isolating the phthalimide deriva-

tive (III), (III) is reacted with an amine represented by general formula (II). Thus, a phthalamide derivative represented by general formula (I-1) can be produced.

5 (1) General formula (V) \rightarrow General formula (III)

As the inert solvent used in this reaction, any solvent may be used so far as it does not disturb the progress of the reaction markedly. Examples of the inert solvent which can be used include aromatic
10 hydrocarbons such as benzene, toluene, xylene and the like, halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride and the like, chlorinated aromatic hydrocarbons such as chlorobenzene, dichlorobenzene and the like, acyclic
15 and cyclic ethers such as diethyl ether, dioxane, tetrahydrofuran and the like, esters such as ethyl acetate and the like, amides such as dimethylformamide, dimethylacetamide and the like, acids such as acetic acid and the like, dimethyl sulfoxide, 1,3-dimethyl-2-
20 imidazolidinone, etc. These inert solvents may be used either singly or in combination or two or more solvents.

Since this reaction is an equimolar reaction, the reactants may be used in equimolar amounts. If
25 desired, however, any one reactant may be also used in excess. According to the need, this reaction may be carried out under a dehydrating condition.

The reaction may be carried out in a tempera-

ture range from room temperature to the refluxing temperature of the used inert solvent. Although the reaction time may vary depending on scale and temperature of the reaction, it may be appropriately selected
5 in a range from several minutes to 48 hours.

After completion of the reaction, the objective product is isolated from the reaction system in the conventional manner and purified by the method of recrystallization, column chromatography, etc.
10 according to the need, whereby the objective product can be obtained. It is also possible to feed the objective product to the subsequent step without isolation.

The phthalic anhydride derivative represented
15 by the general formula (V) can be produced according to the method described in J. Org. Chem., 52, 129 (1987); J. Am. Chem. Soc., 51, 1865 (1929); *ibid.*, 63, 1542 (1941), etc. The heterocyclic amine derivative represented by the general formula (IV) can be produced
20 according to the method described in J. Org. Chem., 18, 138 (1953); J. Org. Chem., 28, 1877 (1963); Chem. Ber., 89, 2742 (1956); Proc. Indian Acad. Sci., 37A, 758 (1953); J. Heterocycl. Chem., 17, 143 (1980); JP-A-62-96479; JP-A-10-340345; JP-A-11-302233; etc.


25 (2) General formula (III) → General formula (I-1)

As the inert solvents usable in this reaction, the same ones as those usable in the above-mentioned reaction (1) can be referred to. Since this

reaction is an equimolar reaction, the reactants may be used in equimolar amounts, though the amine of the general formula (II) may be used in excess, if desired. The reaction can be carried out in a temperature range
5 falling in a range from room temperature to the refluxing temperature of the used inert solvent. Although the reaction time varies depending on scale and temperature of the reaction, it may be selected appropriately in a range from several minutes to 48 hours.

10 After completion of the reaction, the objective product is isolated from the reaction system containing the product in a conventional manner. According to the need, the product is purified by the method of recrystallization, column chromatography,
15 etc., whereby the objective product can be obtained.

Production process 2



wherein R^1 , R^2 , R^3 , X, Q and n are as defined above.

A phthalic anhydride derivative represented by general formula (V) is reacted with an amine represented by general formula (II) in the presence of an inert solvent to form a phthalamide represented by general formula (III-2). In cases where R^1 in (III-2) is hydrogen atom, the phthalamide (III-2) is isolated or not isolated and then subjected to a condensation reaction in the presence of a condensing agent to form a compound represented by general formula (VI-1), and after isolating or without isolating (VI-1), the compound (VI-1) is reacted with a heterocyclic amine derivative represented by general formula (IV) in the presence of an inert solvent. In cases where R^1 in phthalamide (III-2) is not hydrogen atom, (III-2) having been isolated or not isolated is subjected to a condensation reaction with a heterocyclic amine represented by general formula (IV) in the presence of a condensing agent. In these manners, a phthalamide derivative represented by general formula (I-2) can be produced.

Alternatively, a phthalic anhydride derivative represented by general formula (V) is reacted with a heterocyclic amine derivative represented by general formula (IV) in the presence of an inert solvent to form a phthalamide represented by general formula (III-1). In cases where R^3 in (III-1) is hydrogen atom, the phthalamide (III-1) is isolated or not isolated, and

then subjected to a condensation reaction in the presence of a condensing agent to form a compound represented by general formula (VI), and after isolating or not isolating the compound (VI), (VI) is
5 subjected to a reaction with an amine represented by general formula (II) in the presence of an inert solvent. In cases where R³ in phthalamide (III-1) is not hydrogen atom, the phthalamide (III-1) having been isolated or not isolated is subjected to a condensation
10 reaction with an amine represented by general formula (II) in the presence of a condensing agent. In these manners, a phthalamide derivative represented by general formula (I-2) can be obtained.

(1) General formula (V) → General formula (III-2) or
15 General formula (VI-1) → General formula (I-2)

This reaction can be practiced in the same manner as in Production process 1-(2), whereby the objective product can be obtained.

(2) General formula (III-1) → General formula (VI) or
20 General formula (III-2) → General formula (VI-1)

This reaction can be practiced according to the description of J. Med. Chem., 10, 982 (1967), whereby the objective product can be obtained.

(3) General formula (VI) → General formula (I-2) or
25 General formula (V) → General formula (III-2)

This reaction can be practiced in the same manner as in Production process 1-(2), whereby the objective product can be obtained.

(4) General formula (III-1) or General formula (III-2)
→ General formula (I-2)

This reaction can be practiced by reacting a phthalamide derivative represented by general formula
5 (III-1) or (III-2) with an amine represented by general formula (II) or (IV) in the presence of a condensing agent and an inert solvent. This reaction may be practiced in the presence of a base, if necessary.

As examples of the inert solvent used in this
10 reaction, tetrahydrofuran, diethyl ether, dioxane, chloroform, methylene chloride and the like can be referred to. As examples of the condensing agent used in this reaction, those used in the conventional production of amides can be used, of which examples
15 include Mukaiyama reagent (2-chloro-N-methylpyridinium iodide), DCC (1,3-dicyclohexylcarbodiimide), CDI (carbonyl diimidazole), DEPC (diethyl cyanophosphate), etc. The amount of the condensing agent may be appropriately selected in a range from an equimolar
20 amount to an excessive molar amount based on the phthalamide derivative represented by general formula (III-1) or (III-2).

As examples of the base which can be used in this reaction, organic bases such as triethylamine,
25 pyridine and the like, and inorganic bases such as potassium carbonate and the like can be referred to. The amount of the base may be appropriately selected in the range from an equimolar amount to an excessive


molar amount based on the phthalamide derivative represented by general formula (III-1) or (III-2).

The reaction can be carried out in a temperature range from 0°C to the boiling point of the used inert solvent. Although the reaction time may vary depending on scale and temperature of the reaction, it is in the range of from several minutes to 48 hours.

After completion of the reaction, the objective product is isolated from the reaction system by the conventional method, and the product may be purified by recrystallization, column chromatography, etc. according to the need, whereby the objective product can be obtained.

Next, typical examples of the heterocyclic amine derivative represented by general formula (IV') are listed in Table 1, and typical examples of the phthalamide derivative represented by general formula (I) are listed in Tables 2 to 12. The present invention is by no means limited by the compounds shown herein. In the tables shown below, "Me" means methyl, "Et" means ethyl, "Pr" means propyl, "Bu" means butyl, "Ac" means acetyl, "Ph" means phenyl, the expression "c-" means an alicyclic hydrocarbon, "mp" means melting point, and "nD" means refractive index.

General formula (IV'):



(IV')




Table 1

No.	Q'	Ym	mp (°C) or $^1\text{H-NMR}$ [δ (ppm/ CDCl_3)]
IV'-1	Q26	3-Me-5- C_2F_5	2.17 (s. 3H), 4.82 (br. 2H), 7.42 (d. 1H), 8.16 (s. 1H).
IV'-2	Q26	3-Me-5-n- C_3F_7	2.18 (s. 3H), 4.94 (br. 2H), 7.41 (d. 1H), 8.19 (s. 1H).
IV'-3	Q26	3-Me-5-i- C_3F_7	2.18 (s. 3H), 4.80 (br. 2H), 7.42 (d. 1H), 8.15 (s. 1H).

Table 1 (Continued)

No.	Q'	Ym	mp(°C) or ¹ H-NMR [δ (ppm/CDCl ₃)]
IV'-4	Q27	2-n-C ₃ F ₇	4.08 (br. 2H), 7.04 (dd. 1H), 7.43 (d. 1H), 8.16 (d. 1H).
IV'-5	Q27	6-Cl-2-n-C ₃ F ₇	4.65 (br. 2H), 7.17 (d. 1H), 7.57 (d. 1H).
IV'-6	Q27	2-C ₂ F ₅	3.72 (br. 2H), 7.04 (dd. 1H), 7.46 (d. 1H), 8.16 (d. 1H).
IV'-7	Q27	2-i-C ₃ F ₇	4.12 (br. 2H), 7.06 (dd. 1H), 7.44 (dd. 1H), 8.13 (d. 1H).
IV'-8	Q27	4-Me-2-i-C ₃ F ₇	2.22 (s. 3H), 4.12 (br. 2H), 7.34 (d. 1H), 8.07 (s. 1H).
IV'-9	Q27	4-Me-6-i-C ₃ F ₇	2.21 (s. 3H), 4.26 (br. 2H), 7.09 (dd. 1H), 7.98 (d. 1H).
IV'-10	Q27	6-Me-2-i-C ₃ F ₇	2.42 (s. 3H), 4.12 (br. 2H), 6.98 (d. 1H), 7.31 (dd. 1H).
IV'-11	Q27	6-Cl-2-i-C ₃ F ₇	4.40 (br. 2H), 7.12 (d. 1H), 7.41 (dd. 1H).
IV'-12	Q27	6-F-2-i-C ₃ F ₇	
IV'-13	Q27	6-i-C ₃ F ₇	4.28 (br. 2H), 7.04 (dd. 1H), 7.18 (m. 1H), 8.07 (d. 1H).
IV'-14	Q27	4,6-Cl ₂ -2-i-C ₃ F ₇	4.80 (br. 2H), 7.53 (d. 1H).
IV'-15	Q27	6-MeO-2-i-C ₃ F ₇	3.96 (s. 3H), 4.03 (br. 2H), 6.91 (d. 1H), 7.10 (dd. 1H).
IV'-16	Q27	6-MeS-2-i-C ₃ F ₇	2.58 (s. 3H), 4.00 (br. 2H), 6.91 (d. 1H), 7.22 (dd. 1H).
IV'-17	Q27	6-MeSO-2-i-C ₃ F ₇	


Table 1 (Continued)

No.	Q'	Ym	mp(°C) or ¹ H-NMR [δ (ppm/CDCl ₃)]
IV'-18	Q27	6-MeSO ₂ -2-I-C ₃ F ₇	
IV'-19	Q32	4-Me-2-i-C ₃ F ₇	2.46(s. 3H), 3.94(br. 2H), 8.15(s. 1H).
IV'-20	Q32	4-Me-6-i-C ₃ F ₇	2.49(s. 3H), 4.35(br. 2H), 8.55(s. 1H).
IV'-21	Q34	5-i-C ₃ F ₇	5.0(br. 2H), 8.01(s. 1H), 8.31(s. 1H).
IV'-22	Q27	2-OCF ₂ CHF ₂	3.43(br. 2H), 6.13(tt. 1H), 6.88(d. 1H), 7.08(dd. 1H), 7.74(d. 1H).
IV'-23	Q27	2-OCHF ₂	3.60(br. 2H), 6.72(d. 1H), 7.07(dd. 1H), 7.26(dd. 1H), 7.63(d. 1H).
IV'-24	Q27	6-Me-2-OCHF ₂	1.30(s. 3H), 3.45(br. 2H), 6.58(d. 1H), 6.98(d. 1H), 7.30(t. 1H).
IV'-25	Q27	2-SCHF ₂	3.81(br. 2H), 6.94(dd. 1H), 7.24(t. 1H), 7.25(d. 1H), 8.06(d. 1H).
IV'-26	Q27	6-Me-2-SCHF ₂	44-46°C
IV'-27	Q27	2-OCH(CF ₃) ₂	3.70(br. 2H), 6.40(m. 1H), 6.76(d. 1H), 7.08(dd. 1H), 7.59(d. 1H).
IV'-28	Q27	6-Me-2-OCH(CF ₃) ₂	2.33(s. 3H), 3.45(br. 2H), 6.49(m. 1H), 6.64(d. 1H), 7.03(d. 1H).
IV'-29	Q27	6-Cl-2-OCH(CF ₃) ₂	3.89(br. 2H), 6.24(m. 1H), 6.76(d. 1H), 7.16(d. 1H).
IV'-30	Q27	6-F-2-OCH(CF ₃) ₂	
IV'-31	Q27	6-OMe-2-OCH(CF ₃) ₂	3.15-3.60(br. 2H), 3.95(s. 3H), 6.15(m. 1H), 6.38(d. 1H), 6.99(d. 1H).
IV'-32	Q27	6-Cl-2-SCH(CF ₃) ₂	
IV'-33	Q27	6-Me-2-SCH(CF ₃) ₂	

Table 1 (Continued)

No.	Q'	Ym	mp (°C) or ¹ H-NMR [δ (ppm/CDCl ₃)]
IV'-34	Q27	6-F-2-SCH(CF ₃) ₂	
IV'-35	Q27	6-OMe-2-SCH(CF ₃) ₂	
IV'-36	Q27	2-OCF ₂ CHFOCF ₃	
IV'-37	Q27	6-Me-2-OCF ₂ CHFOCF ₃	2.35(s. 3H), 3.50(br. 2H), 6.31(dt. 1H), 6.77(d. 1H), 7.01(d. 1H).
IV'-38	Q27	6-Cl-2-OCF ₂ CHFOCF ₃	
IV'-39	Q27	2-OCF ₂ CHFO-n-C ₃ F ₇	3.20(br. 2H), 6.43(dt. 1H), 6.84(d. 1H), 7.08(dd. 1H), 7.73(d. 1H).
IV'-40	Q27	6-Me-2-OCF ₂ CHFO-n-C ₃ F ₇	2.35(s. 3H), 3.60(br. 2H), 6.50(dt. 1H), 6.74(d. 1H), 7.02(d. 1H).
IV'-41	Q27	6-Cl-2-OCF ₂ CHFO-n-C ₃ F ₇	3.40(br. 2H), 6.37(dt. 1H), 6.85(d. 1H), 7.14(d. 1H).
IV'-42	Q27	6-Me-2-OCF ₂ CHFCF ₃	2.36(s. 3H), 3.30(br. 2H), 5.35(m. 1H), 6.76(d. 1H), 7.01(d. 1H).
IV'-43	Q27	6-Me-2-OCF=CFCF ₃	2.04(s. 3H), 3.10(br. 2H), 6.65(d. 0.5H), 6.69(d. 0.5H), 7.03(d. 1H). (E,Z mixture)
IV'-44	Q27	6-Me-2-OCH(CF ₃) ₂	2.20(s. 3H), 3.20-3.60(br. 2H), 6.41(m. 1H), 6.67(s. 1H), 7.55(s. 1H).
IV'-45	Q27	6-Me-2-OCF ₂ CHF ₂	2.37(s. 3H), 3.40(br. 2H), 6.16(tt. 1H), 6.79(d. 1H), 7.06(d. 1H).
IV'-46	Q27	6-Cl-2-OCF ₂ CHF ₂	3.50(br. 2H), 6.11(tt. 1H), 6.88(d. 1H), 7.15(d. 1H).
IV'-47	Q27	6-Me-2-OCH ₂ C ₂ F ₅	2.31(s. 3H), 3.33(br. 2H), 4.75(t. 2H), 6.55(d. 1H), 6.98(d. 1H).

General formula (I):



Q :





Table 2 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q1	1-1	3-Cl	H	i-Pr	H	O	H	
Q1	1-2	3-Cl	H	i-Pr	H	O	3-Me-5-C ₂ F ₅	
Q1	1-3	3-Cl	H	i-Pr	H	S	4,5-Br ₂	143
Q1	1-4	3-Br	H	i-Pr	H	O	3-Me-5-n-C ₃ F ₇	
Q1	1-5	3-NO ₂	H	i-Pr	H	O	3-Me-5-i-C ₃ F ₇	
Q1	1-6	3-I	H	i-Pr	H	S	H	
Q1	1-7	3-I	H	i-Pr	H	S	3-Me	207
Q1	1-8	3-I	H	i-Pr	H	S	5-Cl	
Q1	1-9	3-I	H	i-Pr	H	S	5-C ₂ F ₅	
Q1	1-10	3-I	H	i-Pr	H	S	5-n-C ₃ F ₇	
Q1	1-11	3-I	H	i-Pr	H	S	5-i-C ₃ F ₇	
Q1	1-12	3-I	H	i-Pr	H	S	3-Me-5-t-Bu	160
Q1	1-13	3-I	H	i-Pr	H	S	3-Me-5-Br	
Q1	1-14	3-I	H	i-Pr	H	S	3-Me-5-C ₂ F ₅	
Q1	1-15	3-I	H	i-Pr	H	S	3-Me-5-n-C ₃ F ₇	
Q1	1-16	3-I	H	i-Pr	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-17	3-I	H	i-Pr	H	S	3-Me-4-C ₂ F ₅	
Q1	1-18	3-I	H	i-Pr	H	S	3-Me-4-n-C ₃ F ₇	
Q1	1-19	3-I	H	i-Pr	H	S	3-Me-4-i-C ₃ F ₇	
Q1	1-20	3-I	H	t-Bu	H	NMe	5-i-C ₃ F ₇	
Q1	1-21	3-I	H	t-Bu	H	NMe	5-C ₂ F ₅	
Q1	1-22	3-I	H	t-Bu	H	NMe	5-n-C ₃ F ₇	
Q1	1-23	3-I	H	t-Bu	H	NMe	3-Me-5-i-C ₃ F ₇	

Table 2 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q1	1-24	3-I	H	CH (CH ₃) CH ₂ SCH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-25	3-I	H	CH (CH ₃) CH ₂ SOCH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-26	3-I	H	CH (CH ₃) CH ₂ SO ₂ CH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-27	3-I	H	C (CH ₃) ₂ CH ₂ SCH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-28	3-I	H	C (CH ₃) ₂ CH ₂ SOCH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-29	3-I	H	C (CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-30	3-I	H	CH (CH ₃) CH ₂ NHAc	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-31	3-I	H	C (CH ₃) ₂ CH ₂ NHAc	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-32	3-I	H	CH (CH ₃) CH ₂ CH ₂ OCH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-33	3-I	H	C (CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-34	3-I	Et	Et	H	O	H	
Q1	1-35	3-I	Et	Et	H	O	3-Me-5-C ₂ F ₅	
Q1	1-36	3-I	Et	Et	H	O	3-Me-5-n-C ₃ F ₇	
Q1	1-37	3-I	Et	Et	H	O	3-Me-5-i-C ₃ F ₇	
Q1	1-38	3-I	Et	Et	H	O	5-Cl	
Q1	1-39	3-I	Et	Et	H	O	5-Br	
Q1	1-40	3-I	Et	Et	H	O	5-n-C ₃ F ₇	
Q1	1-41	6-I	H	i-Pr	H	S	3-Me-5-t-Bu	97
Q1	1-42	6-I	H	i-Pr	H	S	3-Me	168
Q1	1-43	3-CF ₃	H	i-Pr	H	NMe	3-Me-5-C ₂ F ₅	
Q1	1-44	3-Ph	H	i-Pr	H	NMe	3-Me-5-n-C ₃ F ₇	
Q1	1-45	3-SOCF ₃	H	i-Pr	H	NMe	3-Me-5-i-C ₃ F ₇	
Q1	1-46	3-C ₂ F ₅	H	i-Pr	H	NMe	3-Me-5-C ₂ F ₅	

Table 2 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q1	1-47	3-I-4-Cl	H	i-Pr	H	NMe	3-Me-5-n-C ₃ F ₇	
Q1	1-48	3-I-4-CF ₃	H	i-Pr	H	S	3-Me-5-i-C ₃ F ₇	
Q1	1-49	3-CF ₃ -4-Cl	H	i-Pr	H	S	3-Me-5-C ₂ F ₅	
Q1	1-50	3-OCF ₂ O-4	H	i-Pr	H	S	3-Me-5-n-C ₃ F ₇	
Q1	1-51	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	3-Me-5-i-C ₃ F ₇	
Q2	2- 1	3-I	H	i-Pr	H	S	2-Me-5-C ₂ F ₅	
Q2	2- 2	3-I	H	i-Pr	H	S	2-Me-5-n-C ₃ F ₇	
Q2	2- 3	3-I	H	i-Pr	H	S	2-Me-5-i-C ₃ F ₇	
Q2	2- 4	3-I	H	i-Pr	H	S	4-Me-5-C ₂ F ₅	
Q2	2- 5	3-I	H	i-Pr	H	S	4-Me-5-n-C ₃ F ₇	
Q2	2- 6	3-I	H	i-Pr	H	S	4-Me-5-i-C ₃ F ₇	
Q2	2- 7	3-I	H	t-Bu	H	NMe	5-i-C ₃ F ₇	
Q2	2- 8	3-I	H	t-Bu	H	NMe	5-C ₂ F ₅	
Q2	2- 9	3-I	H	t-Bu	H	NMe	5-n-C ₃ F ₇	
Q2	2-10	3-I	H	t-Bu	H	NMe	4-Me-5-i-C ₃ F ₇	
Q2	2-11	3-I	H	CH(CH ₃)CH ₂ SCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q2	2-12	3-I	H	CH(CH ₃)CH ₂ SOCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q2	2-13	3-I	H	CH(CH ₃)CH ₂ SO ₂ CH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q2	2-14	3-I	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q2	2-15	3-I	H	C(CH ₃) ₂ CH ₂ SOCH ₃	H	S	2-Me-5-i-C ₃ F ₇	
Q2	2-16	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	2-Me-5-i-C ₃ F ₇	
Q2	2-17	3-I	H	CH(CH ₃)CH ₂ NHAc	H	S	2-Me-5-i-C ₃ F ₇	
Q2	2-18	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	S	2-Me-5-i-C ₃ F ₇	

Table 2 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q2	2-19	3-I	H	CH(CH ₃)CH ₂ CH ₂ OCH ₃	H	S	2-Me-5-i-C ₃ F ₇	
Q2	2-20	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q2	2-21	3-I	Et	Et	H	O	H	
Q2	2-22	3-I	Et	Et	H	O	2-Me-5-C ₂ F ₅	
Q2	2-23	3-I	Et	Et	H	O	2-Me-5-n-C ₃ F ₇	
Q2	2-24	3-I	Et	Et	H	O	4-Me-5-i-C ₃ F ₇	

Q :






Table 3 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q4	4-1	H	H	i-Pr	H	O	5-Me	185
Q4	4-2	3-Cl	H	i-Pr	H	O	H	
Q4	4-3	3-Cl	H	i-Pr	H	O	4-Me-5-C ₂ F ₅	
Q4	4-4	3-Cl	H	i-Pr	H	O	4,5-Br ₂	
Q4	4-5	3-Cl	H	i-Pr	H	O	5-Me	136
Q4	4-6	3-Cl	H	i-Pr	H	O	5-(4-Br-Ph)	158
Q4	4-7	3-Cl	H	i-Pr	H	O	4-Me-5-(4-Cl-Ph)	184
Q4	4-8	6-Cl	H	i-Pr	H	O	4-Me-5-(4-Cl-Ph)	101
Q4	4-9	3-Br	H	i-Pr	H	O	4-Me-5-n-C ₃ F ₇	
Q4	4-10	3-NO ₂	H	i-Pr	H	O	4-Me-5-i-C ₃ F ₇	
Q4	4-11	3-I	H	i-Pr	H	O	4-Me	144
Q4	4-12	3-I	H	i-Pr	H	O	4-Me-5-CF ₃	151
Q4	4-13	3-I	H	i-Pr	H	S	H	
Q4	4-14	3-I	H	i-Pr	H	S	4-Me	
Q4	4-15	3-I	H	i-Pr	H	S	5-Cl	
Q4	4-16	3-I	H	i-Pr	H	S	5-C ₂ F ₅	
Q4	4-17	3-I	H	i-Pr	H	S	5-n-C ₃ F ₇	
Q4	4-18	3-I	H	i-Pr	H	S	5-i-C ₃ F ₇	
Q4	4-19	3-I	H	i-Pr	H	S	4-Me-5-t-Bu	
Q4	4-20	3-I	H	i-Pr	H	S	4-Me-5-Br	
Q4	4-21	3-I	H	i-Pr	H	S	4-Me-5-C ₂ F ₅	
Q4	4-22	3-I	H	i-Pr	H	S	4-Me-5-n-C ₃ F ₇	
Q4	4-23	3-I	H	i-Pr	H	S	4-Me-5-i-C ₃ F ₇	

Table 3 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q4	4-24	3-I	H	t-Bu	H	NMe	5-i-C ₃ F ₇	
Q4	4-25	3-I	H	t-Bu	H	NMe	5-C ₂ F ₅	
Q4	4-26	3-I	H	t-Bu	H	NMe	5-n-C ₃ F ₇	
Q4	4-27	3-I	H	t-Bu	H	NMe	4-Me-5-i-C ₃ F ₇	
Q4	4-28	3-I	H	CH (CH ₃) CH ₂ SCH	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-29	3-I	H	CH (CH ₃) CH ₂ SOCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-30	3-I	H	CH (CH ₃) CH ₂ SO ₂ CH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-31	3-I	H	C (CH ₃) ₂ CH ₂ SCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-32	3-I	H	C (CH ₃) ₂ CH ₂ SOCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-33	3-I	H	C (CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-34	3-I	H	CH (CH ₃) CH ₂ NHAc	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-35	3-I	H	C (CH ₃) ₂ CH ₂ NHAc	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-36	3-I	H	CH (CH ₃) CH ₂ CH ₂ OCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-37	3-I	H	C (CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-38	3-I	Et	Et	H	O	H	
Q4	4-39	3-I	Et	Et	H	O	4-Me-5-C ₂ F ₅	
Q4	4-40	3-I	Et	Et	H	O	4-Me-5-n-C ₃ F ₇	
Q4	4-41	3-I	Et	Et	H	O	4-Me-5-i-C ₃ F ₇	
Q4	4-42	3-I	Et	Et	H	O	5-Cl	
Q4	4-43	3-I	Et	Et	H	O	5-Br	
Q4	4-44	3-I	Et	Et	H	O	5-n-C ₃ F ₇	
Q4	4-45	6-I	H	i-Pr	H	O	4-Me-5-CF ₃	143
Q4	4-46	3-CF ₃	H	i-Pr	H	NMe	4-Me-5-C ₂ F ₅	

Table 3 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q4	4-47	3-Ph	H	i-Pr	H	NMe	4-Me-5-n-C ₃ F ₇	
Q4	4-48	3-SOCF ₃	H	i-Pr	H	NMe	4-Me-5-i-C ₃ F ₇	
Q4	4-49	3-C ₂ F ₅	H	i-Pr	H	NMe	4-Me-5-C ₂ F ₅	
Q4	4-50	3-I-4-Cl	H	i-Pr	H	NMe	4-Me-5-n-C ₃ F ₇	
Q4	4-51	3-I-4-CF ₃	H	i-Pr	H	S	4-Me-5-i-C ₃ F ₇	
Q4	4-52	3-CF ₃ -4-Cl	H	i-Pr	H	S	4-Me-5-C ₂ F ₅	
Q4	4-53	3-OCF ₂ O-4	H	i-Pr	H	S	4-Me-5-n-C ₃ F ₇	
Q4	4-54	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	4-Me-5-i-C ₃ F ₇	
Q5	5- 1	3-Cl	H	i-Pr	H	O	H	
Q5	5- 2	3-Cl	H	i-Pr	H	O	3-Me-5-Cl	
Q5	5- 3	3-Cl	H	i-Pr	H	O	3,5-Br ₂	
Q5	5- 4	3-Cl	H	i-Pr	H	NMe	3-Me	180
Q5	5- 5	3-Cl	H	i-Pr	H	NMe	3-Me-5-OMe	220
Q5	5- 6	3-Cl	H	n-Pr	H	NMe	3-Me-5-OMe	90
Q5	5- 7	3-Cl	H	n-Pr	H	NMe	3-Me-5-OPh	190
Q5	5- 8	6-Cl	H	i-Pr	H	NMe	3-Me-5-OPh	245
Q5	5- 9	6-Cl	H	i-Pr	H	NMe	3-Me-5-OMe	175
Q5	5-10	3-Br	H	i-Pr	H	O	3,5-Me ₂	
Q5	5-11	3-NO ₂	H	i-Pr	H	O	3,5-Me ₂	
Q5	5-12	3-I	H	i-Pr	H	O	3-CF ₃	
Q5	5-13	3-I	H	i-Pr	H	O	5-CF ₃	
Q5	5-14	3-I	H	i-Pr	H	S	H	
Q5	5-15	3-I	H	i-Pr	H	S	3-Me	

Table 3 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q5	5-16	3-I	H i-Pr		H S		5-Cl	
Q5	5-17	3-I	H i-Pr		H S		5-C ₂ F ₅	
Q5	5-18	3-I	H i-Pr		H S		5-n-C ₃ F ₇	
Q5	5-19	3-I	H i-Pr		H S		5-i-C ₃ F ₇	
Q5	5-20	3-I	H i-Pr		H S		3-C ₂ F ₅	
Q5	5-21	3-I	H i-Pr		H S		3-n-C ₃ F ₇	
Q5	5-22	3-I	H i-Pr		H S		3-i-C ₃ F ₇	
Q5	5-23	3-I	H i-Pr		H S		3-Me-5-Br	
Q5	5-24	3-I	H i-Pr		H S		3-Me-5-C ₂ F ₅	
Q5	5-25	3-I	H i-Pr		H S		3-Me-5-n-C ₃ F ₇	
Q5	5-26	3-I	H i-Pr		H S		3-Me-5-i-C ₃ F ₇	
Q5	5-27	3-I	H t-Bu		H NMe		5-i-C ₃ F ₇	
Q5	5-28	3-I	H t-Bu		H NMe		5-C ₂ F ₅	
Q5	5-29	3-I	H t-Bu		H NMe		5-n-C ₃ F ₇	
Q5	5-30	3-I	H CH (CH ₃) CH ₂ SCH ₃		H NMe		3, 5-Me ₂	
Q5	5-31	3-I	H CH (CH ₃) CH ₂ SOCH ₃		H NMe		3, 5-Me ₂	
Q5	5-32	3-I	H CH (CH ₃) CH ₂ SO ₂ CH ₃		H NMe		3, 5-Me ₂	
Q5	5-33	3-I	H C (CH ₃) ₂ CH ₂ SCH ₃		H NMe		3, 5-Me ₂	
Q5	5-34	3-I	H C (CH ₃) ₂ CH ₂ SOCH ₃		H NMe		3, 5-Me ₂	
Q5	5-35	3-I	H C (CH ₃) ₂ CH ₂ SO ₂ CH ₃		H NMe		3, 5-Me ₂	
Q5	5-36	3-I	H CH (CH ₃) CH ₂ NHAc		H NMe		3, 5-Me ₂	
Q5	5-37	3-I	H C (CH ₃) ₂ CH ₂ NHAc		H NMe		3, 5-Me ₂	
Q5	5-38	3-I	H CH (CH ₃) CH ₂ CH ₂ OCH ₃		H NMe		3, 5-Me ₂	

Table 3 (Continued)


Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q5	5-39	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	NMe	3,5-Me ₂	
Q5	5-40	3-I		Et	Et	H	O	H
Q5	5-41	3-I		Et	Et	H	O	3-Me-5-C ₂ F ₅
Q5	5-42	3-I		Et	Et	H	O	3-Me-5-n-C ₃ F ₇
Q5	5-43	3-I		Et	Et	H	O	3-Me-5-i-C ₃ F ₇
Q5	5-44	3-I		Et	Et	H	O	5-Cl
Q5	5-45	3-I		Et	Et	H	O	5-Br
Q5	5-46	3-I		Et	Et	H	O	5-n-C ₃ F ₇
Q5	5-47	3-I		Et	Et	H	O	5-n-C ₃ F ₇
Q5	5-48	3-CF ₃	H	i-Pr	H	NMe	3-Me-5-C ₂ F ₅	
Q5	5-49	3-Ph	H	i-Pr	H	NMe	3-Me-5-n-C ₃ F ₇	
Q5	5-50	3-SOCF ₃	H	i-Pr	H	NMe	3-Me-5-i-C ₃ F ₇	
Q5	5-51	3-C ₂ F ₅	H	i-Pr	H	NMe	3-Me-5-C ₂ F ₅	
Q5	5-52	3-I-4-Cl	H	i-Pr	H	NMe	3-Me-5-n-C ₃ F ₇	
Q5	5-53	3-I-4-CF ₃	H	i-Pr	H	S	3-Me-5-i-C ₃ F ₇	
Q5	5-54	3-CF ₃ -4-Cl	H	i-Pr	H	S	3-Me-5-C ₂ F ₅	
Q5	5-55	3-OCF ₂ O-4	H	i-Pr	H	S	3-Me-5-n-C ₃ F ₇	
Q5	5-56	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	3-Me-5-i-C ₃ F ₇	
Q6	6- 1	3-Cl	H	i-Pr	H	O	H	
Q6	6- 2	3-Cl	H	i-Pr	H	O	4-Me-3-C ₂ F ₅	
Q6	6- 3	3-Cl	H	i-Pr	H	O	3,4-Br ₂	
Q6	6- 4	3-Br	H	i-Pr	H	O	4-Me-3-n-C ₃ F ₇	
Q6	6- 5	3-NO ₂	H	i-Pr	H	O	4-Me-3-i-C ₃ F ₇	

Table 3 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q6	6- 6	3-NO ₂	H i-Pr		H	NMe	3-Me	176
Q6	6- 7	3-I	H i-Pr		H	O	4-Me-3-Et	85
Q6	6- 8	3-I	H i-Pr		H	O	4-Me-3-CF ₃	103
Q6	6- 9	3-I	H i-Pr		H	S	H	
Q6	6-10	3-I	H i-Pr		H	S	4-Me	
Q6	6-11	3-I	H i-Pr		H	S	3-Cl	
Q6	6-12	3-I	H i-Pr		H	S	3-C ₂ F ₅	
Q6	6-13	3-I	H i-Pr		H	S	3-n-C ₃ F ₇	
Q6	6-14	3-I	H i-Pr		H	S	3-i-C ₃ F ₇	
Q6	6-15	3-I	H i-Pr		H	S	4-Me-3-t-Bu	
Q6	6-16	3-I	H i-Pr		H	S	4-Me-3-Br	
Q6	6-17	3-I	H i-Pr		H	S	4-Me-3-C ₂ F ₅	
Q6	6-18	3-I	H i-Pr		H	S	4-Me-3-n-C ₃ F ₇	
Q6	6-19	3-I	H i-Pr		H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-20	3-I	H t-Bu		H	NMe	3-i-C ₃ F ₇	
Q6	6-21	3-I	H t-Bu		H	NMe	3-C ₂ F ₅	
Q6	6-22	3-I	H t-Bu		H	NMe	3-n-C ₃ F ₇	
Q6	6-23	3-I	H t-Bu		H	NMe	4-Me-3-i-C ₃ F ₇	
Q6	6-24	3-I	H CH(CH ₃)CH ₂ SCH ₃		H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-25	3-I	H CH(CH ₃)CH ₂ SOCH ₃		H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-26	3-I	H CH(CH ₃)CH ₂ SO ₂ CH ₃		H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-27	3-I	H C(CH ₃) ₂ CH ₂ SCH ₃		H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-28	3-I	H C(CH ₃) ₂ CH ₂ SOCH ₃		H	S	4-Me-3-i-C ₃ F ₇	

Table 3 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q6	6-29	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-30	3-I	H	CH(CH ₃)CH ₂ NHAc	H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-31	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-32	3-I	H	CH(CH ₃)CH ₂ CH ₂ OCH ₃	H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-33	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-34	3-I	Et	Et	H	O	H	
Q6	6-35	3-I	Et	Et	H	O	4-Me-3-C ₂ F ₅	
Q6	6-36	3-I	Et	Et	H	O	4-Me-3-n-C ₃ F ₇	
Q6	6-37	3-I	Et	Et	H	O	4-Me-3-i-C ₃ F ₇	
Q6	6-38	3-I	Et	Et	H	O	3-Cl	
Q6	6-39	3-I	Et	Et	H	O	3-Br	
Q6	6-40	3-I	Et	Et	H	O	3-n-C ₃ F ₇	
Q6	6-41	3-CF ₃	H	i-Pr	H	NMe	4-Me-3-C ₂ F ₅	
Q6	6-42	3-Ph	H	i-Pr	H	NMe	4-Me-3-n-C ₃ F ₇	
Q6	6-43	3-SOCF ₃	H	i-Pr	H	NMe	4-Me-3-i-C ₃ F ₇	
Q6	6-44	3-C ₂ F ₅	H	i-Pr	H	NMe	4-Me-3-C ₂ F ₅	
Q6	6-45	3-I-4-Cl	H	i-Pr	H	NMe	4-Me-3-n-C ₃ F ₇	
Q6	6-46	3-I-4-CF ₃	H	i-Pr	H	S	4-Me-3-i-C ₃ F ₇	
Q6	6-47	3-CF ₃ -4-Cl	H	i-Pr	H	S	4-Me-3-C ₂ F ₅	
Q6	6-48	3-OCF ₂ O-4	H	i-Pr	H	S	4-Me-3-n-C ₃ F ₇	
Q6	6-49	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	4-Me-3-i-C ₃ F ₇	

Table 4 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q8	8- 1	3-Cl	H	i-Pr	H	S	H	137
Q8	8- 2	3-Cl	H	i-Pr	H	S	4-Me	175
Q8	8- 3	3-Cl	H	i-Pr	H	S	4-CF ₃	185
Q8	8- 4	3-Cl	H	i-Pr	H	S	4-Ph	175
Q8	8- 5	3-Cl	H	i-Pr	H	S	4-Ph-5-Cl	205
Q8	8- 6	3-Cl	H	i-Pr	H	O	4-Me-5-Cl	
Q8	8- 7	3-Cl	H	i-Pr	H	O	4, 5-Br ₂	
Q8	8- 8	3-Cl	H	i-Pr	H	NMe	4-Me	
Q8	8- 9	3-Cl	H	i-Pr	H	NMe	4-Me-5-OMe	
Q8	8-10	3-Cl	H	n-Pr	H	NMe	4-Me-5-OMe	
Q8	8-11	3-Cl	H	n-Pr	H	NMe	4-Me-5-OPh	
Q8	8-12	6-Cl	H	i-Pr	H	S	4-CH ₃	155
Q8	8-13	6-Cl	H	i-Pr	H	S	4-CF ₃	165
Q8	8-14	6-Cl	H	i-Pr	H	S	4-Ph	155
Q8	8-15	6-Cl	H	i-Pr	H	S	4-Ph-5-Cl	155
Q8	8-16	3-Br	H	i-Pr	H	O	4, 5-Me ₂	
Q8	8-17	3-NO ₂	H	i-Pr	H	O	4, 5-Me ₂	

Table 4 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q8	8-18	3-I	H i-Pr		H	O	4-CF ₃	
Q8	8-19	3-I	H i-Pr		H	O	5-CF ₃	
Q8	8-20	3-I	H i-Pr		H	S	H	
Q8	8-21	3-I	H i-Pr		H	S	4-Me	
Q8	8-22	3-I	H i-Pr		H	S	5-Cl	
Q8	8-23	3-I	H i-Pr		H	S	5-C ₂ F ₅	
Q8	8-24	3-I	H i-Pr		H	S	5-n-C ₃ F ₇	
Q8	8-25	3-I	H i-Pr		H	S	5-i-C ₃ F ₇	
Q8	8-26	3-I	H i-Pr		H	S	4-C ₂ F ₅	
Q8	8-27	3-I	H i-Pr		H	S	4-n-C ₃ F ₇	
Q8	8-28	3-I	H i-Pr		H	S	4-i-C ₃ F ₇	
Q8	8-29	3-I	H i-Pr		H	S	4-Me-5-Br	
Q8	8-30	3-I	H i-Pr		H	S	4-Me-5-C ₂ F ₅	
Q8	8-31	3-I	H i-Pr		H	S	4-Me-5-n-C ₃ F ₇	
Q8	8-32	3-I	H i-Pr		H	S	4-Me-5-i-C ₃ F ₇	
Q8	8-33	3-I	H t-Bu		H	NMe	5-i-C ₃ F ₇	
Q8	8-34	3-I	H t-Bu		H	NMe	5-C ₂ F ₅	
Q8	8-35	3-I	H t-Bu		H	NMe	5-n-C ₃ F ₇	
Q8	8-36	3-I	H CH(CH ₃)CH ₂ SCH ₃		H	NMe	4,5-Me ₂	
Q8	8-37	3-I	H CH(CH ₃)CH ₂ SO ₂ CH ₃		H	NMe	4,5-Me ₂	
Q8	8-38	3-I	H CH(CH ₃)CH ₂ SOCH ₃		H	NMe	4,5-Me ₂	
Q8	8-39	3-I	H C(CH ₃) ₂ CH ₂ SCH ₃		H	NMe	4,5-Me ₂	
Q8	8-40	3-I	H C(CH ₃) ₂ CH ₂ SOCH ₃		H	NMe	4,5-Me ₂	

Table 4 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q8	8-41	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	NMe	4,5-Me ₂	
Q8	8-42	3-I	H	CH(CH ₃)CH ₂ NHAc	H	NMe	4,5-Me ₂	
Q8	8-43	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	NMe	4,5-Me ₂	
Q8	8-44	3-I	H	CH(CH ₃)CH ₂ CH ₂ OCH ₃	H	NMe	4,5-Me ₂	
Q8	8-45	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	NMe	4,5-Me ₂	
Q8	8-46	3-I	Et	Et	H	O	H	
Q8	8-47	3-I	Et	Et	H	O	5-C ₂ F ₅	
Q8	8-48	3-I	Et	Et	H	O	4-n-C ₃ F ₇	
Q8	8-49	3-I	Et	Et	H	O	4-i-C ₃ F ₇	
Q8	8-50	3-I	Et	Et	H	O	5-Cl	
Q8	8-51	3-I	Et	Et	H	O	5-Br	
Q8	8-52	3-I	Et	Et	H	O	5-n-C ₃ F ₇	
Q8	8-53	3-I	Et	Et	H	S	4-(4-Cl-Ph)	139
Q8	8-54	3-CF ₃	H	i-Pr	H	NMe	5-C ₂ F ₅	
Q8	8-55	3-Ph	H	i-Pr	H	NMe	4-n-C ₃ F ₇	
Q8	8-56	3-SOCF ₃	H	i-Pr	H	NMe	4-i-C ₃ F ₇	
Q8	8-57	3-C ₂ F ₅	H	i-Pr	H	NMe	5-C ₂ F ₅	
Q8	8-58	3-I-4-Cl	H	i-Pr	H	NMe	4-n-C ₃ F ₇	
Q8	8-59	3-I-4-CF ₃	H	i-Pr	H	S	4-i-C ₃ F ₇	
Q8	8-60	3-CF ₃ -4-Cl	H	i-Pr	H	S	5-C ₂ F ₅	
Q8	8-61	3-OCF ₂ O-4	H	i-Pr	H	S	4-n-C ₃ F ₇	
Q8	8-62	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	4-i-C ₃ F ₇	
Q8	8-63	3-I	H	i-Pr	H	S	4-S-Et	86

Table 4 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q8	8-64	6-I	H	i-Pr	H	S	4-S-Et	135
Q8	8-65	3-I	H	i-Pr	H	S	4-Me-5-CO ₂ -Et	
							Amorphous solid	
Q9	9- 1	3-Cl	H	i-Pr	H	O	H	
Q9	9- 2	3-Cl	H	i-Pr	H	O	5-Me-2-C ₂ F ₅	
Q9	9- 3	3-Cl	H	i-Pr	H	O	2,5-Br ₂	
Q9	9- 4	3-Cl	H	i-Pr	H	S	2-Ph	131
Q9	9- 5	3-Br	H	i-Pr	H	O	5-Me-2-n-C ₃ F ₇	
Q9	9- 6	3-NO ₂	H	i-Pr	H	O	5-Me-2-i-C ₃ F ₇	
Q9	9- 7	3-I	H	i-Pr	H	O	5-Me-2-CF ₃	
Q9	9- 8	3-I	H	i-Pr	H	S	H	
Q9	9- 9	3-I	H	i-Pr	H	S	2-Me	
Q9	9-10	3-I	H	i-Pr	H	S	2-Cl	
Q9	9-11	3-I	H	i-Pr	H	S	2-C ₂ F ₅	
Q9	9-12	3-I	H	i-Pr	H	S	2-n-C ₃ F ₇	
Q9	9-13	3-I	H	i-Pr	H	S	2-i-C ₃ F ₇	
Q9	9-14	3-I	H	i-Pr	H	S	5-Me-2-t-Bu	
Q9	9-15	3-I	H	i-Pr	H	S	5-Me-2-I	135
Q9	9-16	3-I	H	i-Pr	H	S	5-Me-2-C ₂ F ₅	
Q9	9-17	3-I	H	i-Pr	H	S	5-Me-2-n-C ₃ F ₇	
Q9	9-18	3-I	H	i-Pr	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-19	3-I	H	i-Pr	H	S	5-Me-2-I	191
Q9	9-20	3-I	H	t-Bu	H	NMe	2-i-C ₃ F ₇	

Table 4 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q9	9-21	3-I	H	t-Bu	H	NMe	2-C ₂ F ₅	
Q9	9-22	3-I	H	t-Bu	H	NMe	2-n-C ₃ F ₇	
Q9	9-23	3-I	H	t-Bu	H	NMe	5-Me-2-i-C ₃ F ₇	
Q9	9-24	3-I	H	CH(CH ₃)CH ₂ SCH ₃	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-25	3-I	H	CH(CH ₃)CH ₂ SOCH ₃	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-26	3-I	H	CH(CH ₃)CH ₂ SO ₂ CH ₃	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-27	3-I	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-28	3-I	H	C(CH ₃) ₂ CH ₂ SOCH ₃	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-29	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-30	3-I	H	CH(CH ₃)CH ₂ NHAc	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-31	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-32	3-I	Et	Et	H	O	H	
Q9	9-33	3-I	Et	Et	H	O	5-Me-2-C ₂ F ₅	
Q9	9-34	3-I	Et	Et	H	O	5-Me-2-n-C ₃ F ₇	
Q9	9-35	3-I	Et	Et	H	O	5-Me-2-i-C ₃ F ₇	
Q9	9-36	3-I	Et	Et	H	O	2-Cl	
Q9	9-37	3-I	Et	Et	H	O	2-Br	
Q9	9-38	3-I	Et	Et	H	O	2-n-C ₃ F ₇	
Q9	9-39	3-CF ₃	H	i-Pr	H	NMe	5-Me-2-C ₂ F ₅	
Q9	9-40	3-Ph	H	i-Pr	H	NMe	5-Me-2-n-C ₃ F ₇	
Q9	9-41	3-SOCF ₃	H	i-Pr	H	NMe	5-Me-2-i-C ₃ F ₇	
Q9	9-42	3-C ₂ F ₅	H	i-Pr	H	NMe	5-Me-2-C ₂ F ₅	
Q9	9-43	3-I-4-Cl	H	i-Pr	H	NMe	5-Me-2-n-C ₃ F ₇	

Table 4 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q9	9-44	3-I-4-CF ₃	H	i-Pr	H	S	5-Me-2-i-C ₃ F ₇	
Q9	9-45	3-CF ₃ -4-Cl	H	i-Pr	H	S	5-Me-2-C ₂ F ₅	
Q9	9-46	3-OCF ₂ O-4	H	i-Pr	H	S	5-Me-2-n-C ₃ F ₇	
Q9	9-47	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	5-Me-2-i-C ₃ F ₇	
Q10	10-1	3-Cl	H	i-Pr	H	O	H	
Q10	10-2	3-Cl	H	i-Pr	H	O	4-Me-2-C ₂ F ₅	
Q10	10-3	3-Cl	H	i-Pr	H	O	2,4-Br ₂	
Q10	10-4	3-Cl	H	i-Pr	H	O	2-Ph	
Q10	10-5	3-Br	H	i-Pr	H	O	4-Me-2-n-C ₃ F ₇	
Q10	10-6	3-NO ₂	H	i-Pr	H	O	4-Me-2-i-C ₃ F ₇	
Q10	10-7	3-I	H	i-Pr	H	S	4-Me	230
Q10	10-8	3-I	H	i-Pr	H	O	4-Me-2-CF ₃	
Q10	10-9	3-I	H	i-Pr	H	S	H	
Q10	10-10	3-I	H	i-Pr	H	S	4-Me	
Q10	10-11	3-I	H	i-Pr	H	S	2-Cl	
Q10	10-12	3-I	H	i-Pr	H	S	2-C ₂ F ₅	
Q10	10-13	3-I	H	i-Pr	H	S	2-n-C ₃ F ₇	
Q10	10-14	3-I	H	i-Pr	H	S	2-i-C ₃ F ₇	
Q10	10-15	3-I	H	i-Pr	H	S	4-Me-2-t-Bu	
Q10	10-16	3-I	H	i-Pr	H	S	4-Me-2-I	
Q10	10-17	3-I	H	i-Pr	H	S	4-Me-2-C ₂ F ₅	
Q10	10-18	3-I	H	i-Pr	H	S	4-Me-2-n-C ₃ F ₇	
Q10	10-19	3-I	H	i-Pr	H	S	4-Me-2-i-C ₃ F ₇	

Table 4 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q10	10-20	6-I	H	i-Pr	H	S	4-Me	198
Q10	10-21	3-I	H	t-Bu	H	NMe	2-i-C ₃ F ₇	
Q10	10-22	3-I	H	t-Bu	H	NMe	2-C ₂ F ₅	
Q10	10-23	3-I	H	t-Bu	H	NMe	2-n-C ₃ F ₇	
Q10	10-24	3-I	H	t-Bu	H	NMe	4-Me-2-i-C ₃ F ₇	
Q10	10-25	3-I	H	CH(CH ₃)CH ₂ SCH ₃	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-26	3-I	H	CH(CH ₃)CH ₂ SOCH ₃	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-27	3-I	H	CH(CH ₃)CH ₂ SO ₂ CH ₃	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-28	3-I	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-29	3-I	H	C(CH ₃) ₂ CH ₂ SOCH ₃	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-30	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-31	3-I	H	CH(CH ₃)CH ₂ NHAc	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-32	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-33	3-I	Et	Et	H	O	H	
Q10	10-34	3-I	Et	Et	H	O	4-Me-2-C ₂ F ₅	
Q10	10-35	3-I	Et	Et	H	O	4-Me-2-n-C ₃ F ₇	
Q10	10-36	3-I	Et	Et	H	O	4-Me-2-i-C ₃ F ₇	
Q10	10-37	3-I	Et	Et	H	O	2-Cl	
Q10	10-38	3-I	Et	Et	H	O	2-Br	
Q10	10-39	3-I	Et	Et	H	O	2-n-C ₃ F ₇	
Q10	10-40	3-CF ₃	H	i-Pr	H	NMe	4-Me-2-C ₂ F ₅	
Q10	10-41	3-Ph	H	i-Pr	H	NMe	4-Me-2-n-C ₃ F ₇	
Q10	10-42	3-SOCF ₃	H	i-Pr	H	NMe	4-Me-2-i-C ₃ F ₇	

Table 4 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q10	10-43	3-C ₂ F ₅	H	i-Pr	H	NMe	4-Me-2-C ₂ F ₅	
Q10	10-44	3-I-4-Cl	H	i-Pr	H	NMe	4-Me-2-n-C ₃ F ₇	
Q10	10-45	3-I-4-CF ₃	H	i-Pr	H	S	4-Me-2-i-C ₃ F ₇	
Q10	10-46	3-CF ₃ -4-Cl	H	i-Pr	H	S	4-Me-2-C ₂ F ₅	
Q10	10-47	3-OCF ₂ O-4	H	i-Pr	H	S	4-Me-2-n-C ₃ F ₇	
Q10	10-48	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	4-Me-2-i-C ₃ F ₇	

Q :





Table 5 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q13	13- 1	3-Cl	H	i-Pr	H	S	4-Me	
Q13	13- 2	3-Cl	H	i-Pr	H	O	4-Me	
Q13	13- 3	3-Cl	H	i-Pr	H	NMe	4-Me	
Q13	13- 4	3-I	H	i-Pr	H	S	H	
Q13	13- 5	3-I	H	i-Pr	H	S	4-Me	60
Q13	13- 6	3-I	H	i-Pr	H	S	4-Cl	
Q13	13- 7	3-I	H	i-Pr	H	S	4-CF ₃	
Q13	13- 8	3-I	H	i-Pr	H	S	4-C ₂ F ₅	
Q13	13- 9	3-I	H	i-Pr	H	S	4-n-C ₃ F ₇	
Q13	13-10	3-I	H	i-Pr	H	S	4-i-C ₃ F ₇	
Q13	13-11	3-I	H	i-Pr	H	S	4-t-Bu	
Q13	13-12	6-I	H	i-Pr	H	S	4-Me	73
Q13	13-13	3-I	H	CH (CH ₃) CH ₂ SCH ₃	H	S	4-CF ₃	
Q13	13-14	3-I	H	CH (CH ₃) CH ₂ SOCH ₃	H	S	4-CF ₃	
Q13	13-15	3-I	H	CH (CH ₃) CH ₂ SO ₂ CH ₃	H	S	4-CF ₃	
Q13	13-16	3-I	H	C (CH ₃) ₂ CH ₂ SCH ₃	H	S	4-CF ₃	
Q13	13-17	3-I	H	C (CH ₃) ₂ CH ₂ SOCH ₃	H	S	4-CF ₃	
Q13	13-18	3-I	H	C (CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	4-CF ₃	
Q13	13-19	3-I	H	CH (CH ₃) CH ₂ NHAc	H	S	4-CF ₃	
Q13	13-20	3-I	H	C (CH ₃) ₂ CH ₂ NHAc	H	S	4-CF ₃	
Q13	13-21	3-I	H	CH (CH ₃) CH ₂ CH ₂ OCH ₃	H	S	4-CF ₃	
Q13	13-22	3-I	H	C (CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	S	4-CF ₃	
Q13	13-23	3-I	Et	Et	H	S	H	

Table 5 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q13	13-24	3-I	Et	Et	H	S	4-CF ₃	
Q13	13-25	3-I	Et	Et	H	S	4-CF ₃	
Q13	13-26	3-I	Et	Et	H	S	4-CF ₃	
Q13	13-27	3-I	Et	Et	H	S	4-CF ₃	
Q13	13-28	3-CF ₃	H	i-Pr	H	S	3-C ₂ F ₅	
Q13	13-29	3-Ph	H	i-Pr	H	S	3-n-C ₃ F ₇	
Q13	13-30	3-SOCF ₃	H	i-Pr	H	S	3-i-C ₃ F ₇	
Q13	13-31	3-C ₂ F ₅	H	i-Pr	H	S	3-C ₂ F ₅	
Q13	13-32	3-I-4-Cl	H	i-Pr	H	S	3-n-C ₃ F ₇	
Q13	13-33	3-I-4-CF ₃	H	i-Pr	H	S	3-i-C ₃ F ₇	
Q13	13-34	3-CF ₃ -4-Cl	H	i-Pr	H	S	3-C ₂ F ₅	
Q13	13-35	3-OCF ₂ O-4	H	i-Pr	H	S	3-n-C ₃ F ₇	
Q13	13-36	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	3-i-C ₃ F ₇	

Q :




Q19

Table 6 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q19	19- 1	3-Cl	H i-Pr		H	O	H	
Q19	19- 2	3-Cl	H i-Pr		H	O	5-C ₂ F ₅	
Q19	19- 3	3-Cl	H i-Pr		H	S	5-Me	166
Q19	19- 4	3-Br	H i-Pr		H	O	5-n-C ₃ F ₇	
Q19	19- 5	3-NO ₂	H i-Pr		H	O	5-i-C ₃ F ₇	
Q19	19- 6	3-I	H i-Pr		H	S	H	
Q19	19- 7	3-I	H i-Pr		H	S	5-Me	
Q19	19- 8	3-I	H i-Pr		H	S	5-Cl	
Q19	19- 9	3-I	H i-Pr		H	S	5-CF ₃	104
Q19	19-10	3-I	H i-Pr		H	S	5-C ₂ F ₅	
Q19	19-11	3-I	H i-Pr		H	S	5-n-C ₃ F ₇	
Q19	19-12	3-I	H i-Pr		H	S	5-i-C ₃ F ₇	
Q19	19-13	3-I	H i-Pr		H	S	5-t-Bu	
Q19	19-14	3-I	H i-Pr		H	S	5-CF ₃	176
Q19	19-15	3-I	H t-Bu		H	NMe	5-i-C ₃ F ₇	
Q19	19-16	3-I	H t-Bu		H	NMe	5-C ₂ F ₅	
Q19	19-17	3-I	H t-Bu		H	NMe	5-n-C ₃ F ₇	
Q19	19-18	3-I	H t-Bu		H	NMe	5-CF ₃	
Q19	19-19	3-I	H CH(CH ₃)CH ₂ SCH ₃		H	S	5-i-C ₃ F ₇	
Q19	19-20	3-I	H CH(CH ₃)CH ₂ SOCH ₃		H	S	5-i-C ₃ F ₇	
Q19	19-21	3-I	H CH(CH ₃)CH ₂ SO ₂ CH ₃		H	S	5-i-C ₃ F ₇	
Q19	19-22	3-I	H C(CH ₃) ₂ CH ₂ SCH ₃		H	S	5-i-C ₃ F ₇	
Q19	19-23	3-I	H C(CH ₃) ₂ CH ₂ SOCH ₃		H	S	5-i-C ₃ F ₇	

Table 6 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	mp (°C)
Q19	19-24	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	S	5-i-C ₃ F ₇	
Q19	19-25	3-I	H	CH(CH ₃)CH ₂ NHAc	H	S	5-i-C ₃ F ₇	
Q19	19-26	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	S	5-i-C ₃ F ₇	
Q19	19-27	3-I	H	CH(CH ₃)CH ₂ CH ₂ OCH ₃	H	S	5-i-C ₃ F ₇	
Q19	19-28	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	S	5-i-C ₃ F ₇	
Q19	19-29	3-I	Et	Et	H	O	H	
Q19	19-30	3-I	Et	Et	H	O	5-C ₂ F ₅	
Q19	19-31	3-I	Et	Et	H	O	5-n-C ₃ F ₇	
Q19	19-32	3-I	Et	Et	H	O	5-i-C ₃ F ₇	
Q19	19-33	3-I	Et	Et	H	O	5-Cl	
Q19	19-34	3-I	Et	Et	H	S	5-t-Bu	59
Q19	19-35	3-CF ₃	H	i-Pr	H	NMe	5-C ₂ F ₅	
Q19	19-36	3-Ph	H	i-Pr	H	NMe	5-n-C ₃ F ₇	
Q19	19-37	3-SOCF ₃	H	i-Pr	H	NMe	5-i-C ₃ F ₇	
Q19	19-38	3-C ₂ F ₅	H	i-Pr	H	NMe	5-C ₂ F ₅	
Q19	19-39	3-I-4-Cl	H	i-Pr	H	NMe	5-n-C ₃ F ₇	
Q19	19-40	3-I-4-CF ₃	H	i-Pr	H	S	5-i-C ₃ F ₇	
Q19	19-41	3-CF ₃ -4-Cl	H	i-Pr	H	S	5-C ₂ F ₅	
Q19	19-42	3-OCF ₂ O-4	H	i-Pr	H	S	5-n-C ₃ F ₇	
Q19	19-43	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	S	5-i-C ₃ F ₇	

Table 7 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q26	26- 1	H	H	i-Pr	H	3-Cl-5-CF ₃	85
Q26	26- 2	3-Cl	H	i-Pr	H	H	
Q26	26- 3	3-Cl	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q26	26- 4	3-Br	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q26	26- 5	3-NO ₂	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	
Q26	26- 6	3-I	H	i-Pr	H	5-C ₂ F ₅	
Q26	26- 7	3-I	H	i-Pr	H	5-n-C ₃ F ₇	
Q26	26- 8	3-I	H	i-Pr	H	5-i-C ₃ F ₇	
Q26	26- 9	3-I	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q26	26-10	3-I	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q26	26-11	3-I	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	140
Q26	26-12	3-I	H	i-Pr	H	3-Me-4-C ₂ F ₅	
Q26	26-13	3-I	H	i-Pr	H	3-Me-4-n-C ₃ F ₇	
Q26	26-14	3-I	H	i-Pr	H	3-Me-4-i-C ₃ F ₇	
Q26	26-15	3-I	H	t-Bu	H	5-i-C ₃ F ₇	
Q26	26-16	3-I	H	t-Bu	H	5-C ₂ F ₅	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q26	26-17	3-I	H	t-Bu	H	5-n-C ₃ F ₇	
Q26	26-18	3-I	H	t-Bu	H	3-Me-5-i-C ₃ F ₇	
Q26	26-19	3-I	H	CH(CH ₃)CH ₂ SCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-20	3-I	H	CH(CH ₃)CH ₂ SOCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-21	3-I	H	CH(CH ₃)CH ₂ SO ₂ CH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-22	3-I	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-23	3-I	H	C(CH ₃) ₂ CH ₂ SOCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-24	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-25	3-I	H	CH(CH ₃)CH ₂ SEt	H	3-Me-5-i-C ₃ F ₇	
Q26	26-26	3-I	H	C(CH ₃)CH ₂ SEt	H	3-Me-5-i-C ₃ F ₇	
Q26	26-27	3-I	H	CH(CH ₃)CH ₂ CH ₂ SCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-28	3-I	H	CH(CH ₃) ₂ CH ₂ CH ₂ SCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q26	26-29	3-I	Et	Et	H	3-Me-5-C ₂ F ₅	Paste
Q26	26-30	3-I	Et	Et	H	3-Me-5-n-C ₃ F ₇	Paste
Q26	26-31	3-I	Et	Et	H	3-Me-5-i-C ₃ F ₇	159
Q26	26-32	3-I	Et	Et	H	5-Cl	127
Q26	26-33	3-I	Et	Et	H	5-Br	154
Q26	26-34	3-CF ₃	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q26	26-35	3-Ph	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q26	26-36	3-SOCF ₃	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	
Q26	26-37	3-C ₂ F ₅	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q26	26-38	3-I-4-Cl	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q26	26-39	3-I-4-CF ₃	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q26	26-40	3-CF ₃ -4-Cl	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q26	26-41	3-OCF ₂ O-4	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q26	26-42	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	
Q26	26-43	3-I	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	140
Q27	27- 1	H	H	i-Pr	H	H	139
Q27	27- 2	H	H	i-Pr	H	2-Me	
Q27	27- 3	H	H	i-Pr	H	3-Me	
Q27	27- 4	H	H	i-Pr	H	4-Me	
Q27	27- 5	H	H	i-Pr	H	6-Me	
Q27	27- 6	H	H	i-Pr	H	2-Cl	
Q27	27- 7	H	H	i-Pr	H	3-Cl	
Q27	27- 8	H	H	i-Pr	H	4-Cl	
Q27	27- 9	H	H	i-Pr	H	6-Cl	
Q27	27-10	3-Cl	H	i-Pr	H	2-CF ₃	
Q27	27-11	3-Cl	H	i-Pr	H	3-CF ₃	
Q27	27-12	3-Cl	H	i-Pr	H	4-CF ₃	
Q27	27-13	3-Cl	H	i-Pr	H	6-CF ₃	
Q27	27-14	3-Cl	H	i-Pr	H	2-NO ₂	
Q27	27-15	3-Cl	H	i-Pr	H	3-NO ₂	
Q27	27-16	3-Cl	H	i-Pr	H	4-NO ₂	
Q27	27-17	3-Cl	H	i-Pr	H	6-NO ₂	
Q27	27-18	3-Cl	H	i-Pr	H	2-Et	
Q27	27-19	3-Cl	H	i-Pr	H	2-i-Pr	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-20	3-Cl	H	i-Pr	H	2-t-Bu	
Q27	27-21	3-Cl	H	i-Pr	H	2-SCH ₃	
Q27	27-22	3-Cl	H	i-Pr	H	2-SOCH ₃	
Q27	27-23	3-Cl	H	i-Pr	H	2-SO ₂ CH ₃	
Q27	27-24	3-Cl	H	i-Pr	H	2-SCF ₃	
Q27	27-25	3-Cl	H	i-Pr	H	2-SCHF ₂	
Q27	27-26	3-Cl	H	i-Pr	H	2-COCH ₃	
Q27	27-27	3-Cl	H	i-Pr	H	2-CN	
Q27	27-28	3-Cl	H	i-Pr	H	2-OCH ₃	
Q27	27-29	3-Cl	H	i-Pr	H	2-O- (4-Br-Ph)	101
Q27	27-30	3-Cl	H	i-Pr	H	2-O- (2, 4-Cl ₂ -Ph)	97
Q27	27-31	3-Cl	H	i-Pr	H	4-S-i-Pr	193
Q27	27-32	3-Cl	H	i-Pr	H	4-S-i-Bu	183
Q27	27-33	3-Cl	H	i-Pr	H	2-OCF ₂ CCl ₂ F	
Q27	27-34	3-Cl	H	i-Pr	H	2-OCH ₂ CF ₃	
Q27	27-35	3-Cl	H	i-Pr	H	2-OCH ₂ CF ₂ CHF ₂	
Q27	27-36	3-Cl	H	i-Pr	H	2-CF ₂ CF ₃	
Q27	27-37	3-Cl	H	i-Pr	H	2-COOCH (CF ₃) ₂	
Q27	27-38	3-Cl	H	i-Pr	H	2-Ph	
Q27	27-39	3-Cl	H	n-Bu	H	2-Cl	
Q27	27-40	3-Cl	H	i-Bu	H	2-Cl	
Q27	27-41	3-Cl	H	s-Bu	H	2-Cl	
Q27	27-42	3-Cl	H	t-Bu	H	2-Cl	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-43	3-Cl	H	C-C ₃ H ₅	H	2-Cl	
Q27	27-44	3-Cl	H	C-C ₄ H ₇	H	2-Cl	
Q27	27-45	3-Cl	H	C-C ₅ H ₉	H	2-Cl	
Q27	27-46	3-Cl	H	C-C ₆ H ₁₁	H	2-Cl	
Q27	27-47	3-Cl	H	CH ₂ CH=CH ₂	H	2-Cl	
Q27	27-48	3-Cl	H	CH ₂ C≡CH	H	2-Cl	
Q27	27-49	3-Cl	H	CH ₂ Ph	H	2-Cl	
Q27	27-50	3-Cl	H	C(CH ₃) ₂ C≡CH	H	2-Cl	
Q27	27-51	3-Cl	H	C(CH ₃) ₂ C≡CPh	H	2-Cl	
Q27	27-52	3-Cl	H	CH ₂ CH ₂ SCH ₃	H	2-Cl	
Q27	27-53	3-Cl	H	CH ₂ CH ₂ SPh	H	2-Cl	
Q27	27-54	3-Cl	H	CH ₂ CH ₂ SO ₂ Ph	H	2-Cl	
Q27	27-55	3-Cl	H	CH ₂ CH ₂ SO ₂ CH ₃	H	2-Cl	
Q27	27-56	3-Cl	H	CH ₂ CH ₂ CO ₂ CH ₃	H	2-Cl	
Q27	27-57	3-Cl	H	CH ₂ CH ₂ CONHCH ₃	H	2-Cl	
Q27	27-58	3-Cl	Et	Et	H	2-Cl	
Q27	27-59	3-Cl	n-Pr	n-Pr	H	2-Cl	
Q27	27-60	3-Cl	i-Pr	i-Pr	H	2-Cl	
Q27	27-61	3-Cl	i-Pr	Me	H	2-Cl	
Q27	27-62	3-Cl	i-Bu	Me	H	2-Cl	
Q27	27-63	3-Cl	CH ₂ CH=CH ₂	CH ₂ CH=CH ₂	H	2-Cl	
Q27	27-64	3-Cl	Et	Et	Me	2-Cl	
Q27	27-65	3-Cl	n-Pr	i-Pr	Me	2-Cl	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-66	3-Cl	i-Pr	i-Pr	Me	2-Cl	
Q27	27-67	3-Cl	Et	Et	Ac	2-Cl	
Q27	27-68	3-Cl	n-Pr	i-Pr	Ac	2-Cl	
Q27	27-69	3-Cl	i-Pr	i-Pr	Ac	2-Cl	
Q27	27-70	3-Cl		-(CH ₂) ₄ -	H	2-Cl	
Q27	27-71	3-Cl		-(CH ₂) ₂ O(CH ₂) ₂ -	H	2-Cl	
Q27	27-72	3-Cl	i-Pr	SO ₂ CH ₃	H	2-Cl	
Q27	27-73	3-Cl	i-Pr	CN	H	2-Cl	
Q27	27-74	3-Cl	i-Pr	CO ₂ CH ₃	H	2-Cl	
Q27	27-75	3-Cl	i-Pr	COCH ₃	H	2-Cl	
Q27	27-76	3-Cl	i-Pr	COPh	H	2-Cl	
Q27	27-77	3-Cl	i-Pr	NHCOCH ₃	H	2-Cl	
Q27	27-78	3-Cl	H	i-Pr	H	2,4-Me ₂	
Q27	27-79	3-Cl	H	i-Pr	H	2,4-Cl ₂	
Q27	27-80	3-Cl	H	i-Pr	H	4,6-Me ₂	
Q27	27-81	3-Cl	H	i-Pr	H	4-Me-2-Cl	211
Q27	27-82	3-Cl	H	i-Pr	H	4-Me-2-F	
Q27	27-83	3-Cl	H	i-Pr	H	4-Me-2-Br	
Q27	27-84	3-Cl	H	i-Pr	H	4-Me-2-I	
Q27	27-85	3-Cl	H	i-Pr	H	4-Me-2-OCHF ₂	
Q27	27-86	3-Cl	H	i-Pr	H	4-Me-2-OCF ₃	
Q27	27-87	3-Cl	H	i-Pr	H	4-Me-2-NO ₂	
Q27	27-88	3-Cl	H	i-Pr	H	4-Me-2-NMe ₂	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-89	3-Cl	H	i-Pr	H	4-Me-2-C≡CH	
Q27	27-90	3-Cl	H	i-Pr	H	4-Me-2-C≡C-t-Bu	
Q27	27-91	3-Cl	H	i-Pr	H	4-Me-2-C≡CPh	
Q27	27-92	3-Cl	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-93	3-Cl	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-94	3-Cl	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-95	3-Cl	H	i-Pr	H	4-Me-2-OCH ₂ OCH ₃	
Q27	27-96	3-Cl	H	i-Pr	H	4-Me-2-OCF ₂ CHF ₂	
Q27	27-97	3-Cl	H	i-Pr	H	4-Me-2-OPh	
Q27	27-98	3-Cl	H	i-Pr	H	4-Me-2-O-(4-Br-Ph)	79
Q27	27-99	3-Cl	H	i-Pr	H	4-Me-2-OSO ₂ Ph	
Q27	27-100	3-Cl	H	i-Pr	H	4-Me-2-OCH ₂ CO ₂ CH ₃	
Q27	27-101	3-Cl	H	i-Pr	H	4-Me-2-CO ₂ CH ₃	
Q27	27-102	3-Cl	H	i-Pr	H	4-Me-2-S-i-Pr	
Q27	27-103	3-Cl	H	i-Pr	H	4-Me-2-SCHF ₂	
Q27	27-104	3-Cl	H	i-Pr	H	4-Me-2-SOCHF ₂	
Q27	27-105	3-Cl	H	i-Pr	H	4-Me-2-SO ₂ CHF ₂	
Q27	27-106	3-Cl	H	i-Pr	H	4-Cl-2-CF ₃	
Q27	27-107	3-Cl	H	i-Pr	H	4-Cl-2-OCF ₃	
Q27	27-108	3-Cl	H	i-Pr	H	4-Cl-2-i-C ₃ F ₇	
Q27	27-109	3-Cl	H	i-Pr	H	4-Cl-2-C ₂ F ₅	
Q27	27-110	3-Cl	H	i-Pr	H	4-Cl-2-OCHF ₂	
Q27	27-111	3-Cl	H	i-Pr	H	4-Cl-2-OSO ₂ Ph	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-112	3-Cl	H	i-Pr	H	4-OCH ₃ -2-Ph	
Q27	27-113	3-Cl	H	i-Pr	H	4-CF ₃ -2-Cl	
Q27	27-114	3-Cl	H	i-Pr	H	4-Me-3-CF ₃	
Q27	27-115	3-Cl	H	i-Pr	H	4-Me-3-Cl	
Q27	27-116	3-Cl	H	i-Pr	H	4-Me-3-OCF ₃	
Q27	27-117	3-Cl	H	i-Pr	H	4-Me-3-CF ₂ CF ₃	
Q27	27-118	3-Cl	H	i-Pr	H	4-Me-3-n-C ₃ F ₇	
Q27	27-119	3-Cl	H	i-Pr	H	4-Me-3-i-C ₃ F ₇	
Q27	27-120	3-Cl	H	i-Pr	H	3,4-Me ₂ -2-Cl	
Q27	27-121	3-Cl	H	i-Pr	H	3,4-Me ₂ -2-OMe	
Q27	27-122	3-Cl	H	i-Pr	H	3,4-Me ₂ -2-SMe	
Q27	27-123	3-Cl	H	i-Pr	H	4-Me-2,3-Cl ₂	
Q27	27-124	6-Cl	H	i-Pr	H	2-O- (4-Br-Ph)	170
Q27	27-125	6-Cl	H	i-Pr	H	2-O- (2,4-Cl ₂ -Ph)	189
Q27	27-126	6-Cl	H	i-Pr	H	2-S-i-Pr	120
Q27	27-127	6-Cl	H	i-Pr	H	2-S-i-Bu	187
Q27	27-128	6-Cl	H	i-Pr	H	4-Me-2-Cl	230
Q27	27-129	3-I	Et	Et	H	6-Cl-2-n-C ₃ F ₇	122
Q27	27-130	3-I	Et	Et	H	2-Cl	203
Q27	27-131	3-I	Et	Et	H	2-n-C ₃ F ₇	200
Q27	27-132	3-I	Et	Et	H	2-O- (4-Br-Ph)	247
Q27	27-133	3-I	H	i-Pr	H	2-Cl	215
Q27	27-134	3-I	H	i-Pr	H	2-C ₂ F ₅	Amorphous solid

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-135	3-I	H i-Pr		H	2-n-C ₃ F ₇	200
Q27	27-136	3-I	H i-Pr		H	2-i-C ₃ F ₇	270
Q27	27-137	3-I	H i-Pr		H	4-Me-2-i-C ₃ F ₇	257
Q27	27-138	3-I	H i-Pr		H	6-Me-2-i-C ₃ F ₇	234
Q27	27-139	3-I	H t-Bu		H	2-i-C ₃ F ₇	275
Q27	27-140	3-I	H t-Bu		H	2-C ₂ F ₅	260
Q27	27-141	3-I	H t-Bu		H	2-n-C ₃ F ₇	245
Q27	27-142	3-I	H t-Bu		H	4-Me-2-i-C ₃ F ₇	250
Q27	27-143	3-I	H t-Bu		H	6-Me-2-i-C ₃ F ₇	246
Q27	27-144	3-I	H CH (CH ₃) CH ₂ SCH ₃		H	4-Me-2-i-C ₃ F ₇	225
Q27	27-145	3-I	H CH (CH ₃) CH ₂ SCH ₃		H	6-Me-2-i-C ₃ F ₇	229
Q27	27-146	3-I	H CH (CH ₃) CH ₂ SOCH ₃		H	4-Me-2-i-C ₃ F ₇	
Q27	27-147	3-I	H CH (CH ₃) CH ₂ SOCH ₃		H	6-Me-2-i-C ₃ F ₇	
Q27	27-148	3-I	H CH (CH ₃) CH ₂ SO ₂ CH ₃		H	4-Me-2-i-C ₃ F ₇	
Q27	27-149	3-I	H CH (CH ₃) CH ₂ SO ₂ CH ₃		H	6-Me-2-i-C ₃ F ₇	
Q27	27-150	3-I	H C (CH ₃) ₂ CH ₂ SCH ₃		H	4-Me-2-i-C ₃ F ₇	173
Q27	27-151	3-I	H C (CH ₃) ₂ CH ₂ SCH ₃		H	6-Me-2-i-C ₃ F ₇	213
Q27	27-152	3-I	H C (CH ₃) ₂ CH ₂ SOCH ₃		H	4-Me-2-i-C ₃ F ₇	
Q27	27-153	3-I	H C (CH ₃) ₂ CH ₂ SOCH ₃		H	6-Me-2-i-C ₃ F ₇	Amorphous
Q27	27-154	3-I	H C (CH ₃) ₂ CH ₂ SO ₂ CH ₃		H	4-Me-2-i-C ₃ F ₇	
Q27	27-155	3-I	H C (CH ₃) ₂ CH ₂ SO ₂ CH ₃		H	6-Me-2-i-C ₃ F ₇	
Q27	27-156	3-I	H i-Pr		H	4-Me-2-i-C ₃ F ₇	
Q27	27-157	3-I	H t-Bu		H	6-Me-2-i-C ₃ F ₇	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-158	3-F	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	4-Me-2-i-C ₃ F ₇	
Q27	27-159	3-F	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	6-Me-2-i-C ₃ F ₇	
Q27	27-160	3-Br	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-161	3-Br	H	t-Bu	H	6-Me-2-i-C ₃ F ₇	
Q27	27-162	3-Br	H	CH(CH ₃)CH ₂ SCH ₃	H	4-Me-2-i-C ₃ F ₇	
Q27	27-163	3-Br	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	6-Me-2-i-C ₃ F ₇	
Q27	27-164	3-NO ₂	H	i-Pr	H	H	209
Q27	27-165	3-NO ₂	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-166	3-NO ₂	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-167	3-NO ₂	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-168	3-NO ₂	H	i-Pr	H	6-Me-2-CF ₂ CF ₃	
Q27	27-169	3-NO ₂	H	i-Pr	H	6-Me-2-n-C ₃ F ₇	
Q27	27-170	3-NO ₂	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-171	3-NO ₂	H	i-Pr	H	4-Me-2-Cl	
Q27	27-172	3-CN	Et	Et	H	4-Me-2-CF ₂ CF ₃	
Q27	27-173	3-CN	Et	Et	H	4-Me-2-n-C ₃ F ₇	
Q27	27-174	3-CN	Et	Et	H	4-Me-2-i-C ₃ F ₇	
Q27	27-175	3-CN	Et	Et	H	6-Me-2-CF ₂ CF ₃	
Q27	27-176	3-CN	Et	Et	H	6-Me-2-n-C ₃ F ₇	
Q27	27-177	3-CN	Et	Et	H	6-Me-2-i-C ₃ F ₇	
Q27	27-178	3-CN	Et	Et	H	4-Me-2-Cl	
Q27	27-179	3-CF ₃	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-180	3-CF ₃	H	i-Pr	H	6-Me-2-n-C ₃ F ₇	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-181	3-OCH ₃	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-182	3-OCH ₃	H	i-Pr	H	6-Me-2-CF ₂ CF ₃	
Q27	27-183	3-OCH ₃	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-184	3-OCH ₃	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-185	3-SCH ₃	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-186	3-SCH ₃	H	i-Pr	H	6-Me-2-n-C ₃ F ₇	
Q27	27-187	3-S-i-Pr	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-188	3-S-i-Pr	H	i-Pr	H	6-Me-2-CF ₂ CF ₃	
Q27	27-189	3-SOCH ₃	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-190	3-SOCH ₃	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-191	3-SO ₂ CH ₃	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-192	3-SO ₂ CH ₃	H	i-Pr	H	6-Me-2-n-C ₃ F ₇	
Q27	27-193	3-SCH ₂ CF ₃	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-194	3-SCF ₃	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-195	3-SOCF ₃	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-196	3-SO ₂ CF ₃	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-197	3-SPh	H	i-Pr	H	6-Me-2-CF ₂ CF ₃	
Q27	27-198	3-SOPh	H	i-Pr	H	6-Me-2-n-C ₃ F ₇	
Q27	27-199	3-SO ₂ Ph	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-200	3-OPh	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-201	3-Ph	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-202	3-C≡CH	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-203	3-C≡C-t-Bu	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-204	3-C≡CPh	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-205	3-C ₂ F ₅	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-206	3-CO ₂ CH ₃	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-207	3-CONHCH ₃	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-208	3-COCH ₃	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-209	3-CCH ₃ (=NOCH ₃)	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-210	3,4-Cl ₂	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-211	3,6-Cl ₂	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-212	3,5-Cl ₂	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-213	3,5-Cl ₂	H	i-Pr	H	6-Me-2-CF ₂ CF ₃	
Q27	27-214	4,5-Cl ₂	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-215	4,5-Cl ₂	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-216	3-I-4-Cl	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-217	3-I-4-F	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-218	3-I-4CF ₃	H	i-Pr	H	4-Me-2-i-C ₃ F ₇	
Q27	27-219	3-I-4-OCH ₃	H	i-Pr	H	4-Me-2-CF ₂ CF ₃	
Q27	27-220	3-CF ₃ -4-Cl	H	i-Pr	H	4-Me-2-n-C ₃ F ₇	
Q27	27-221	3-CF ₃ -4-OCH ₃	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-222	3-OCH ₂ O-4	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-223	3-OCF ₂ O-4	H	i-Pr	H	6-Me-2-CF ₂ CF ₃	
Q27	27-224	3-OCH ₂ CH ₂ O-4	H	i-Pr	H	6-Me-2-n-C ₃ F ₇	
Q27	27-225	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	
Q27	27-226	3-CH=CH-CH=CH-4	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-227	3-I	H		i-Pr	H 4-Me-3-CF ₂ CF ₃	
Q27	27-228	3-I	H		i-Pr	H 4-Me-3-i-C ₃ F ₇	
Q27	27-229	3-I	H		i-Pr	H 4-Me-3-n-C ₃ F ₇	
Q27	27-230	3-I	H		i-Pr	H 6-Cl-2-i-C ₃ F ₇	188
Q27	27-231	3-I	Et		Et	H 6-Cl-2-i-C ₃ F ₇	164
Q27	27-232	3-I	H	CH(Me)CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	177
Q27	27-233	3-I	H	C(Me) ₂ CH ₂ SMe	H	2-i-C ₃ F ₇	229
Q27	27-234	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	175
Q27	27-235	3-I	H	C(Me) ₂ CH ₂ SOMe	H	6-Cl-2-i-C ₃ F ₇	Paste
Q27	27-236	3-I	H	C(Me) ₂ CH ₂ SO ₂ Me	H	6-Cl-2-i-C ₃ F ₇	Amorphous
Q27	27-237	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-i-C ₃ F ₇	183
Q27	27-238	3-I	H	C(Me) ₂ CH ₂ SOMe	H	6-i-C ₃ F ₇	Amorphous
Q27	27-239	3-I	H	C(Me) ₂ CH ₂ SO ₂ Me	H	6-i-C ₃ F ₇	Amorphous
Q27	27-240	3-I	H	C(Me) ₂ CH ₂ SMe	H	4,6-Cl ₂ -2-i-C ₃ F ₇	120
Q27	27-241	3-I	H	CH(Me)CH ₂ SMe	H	6-MeO-2-i-C ₃ F ₇	134
Q27	27-242	3-I	H		i-Pr	H 6-MeO-2-i-C ₃ F ₇	158
Q27	27-243	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-MeO-2-i-C ₃ F ₇	134
Q27	27-244	3-I	H	C(Me) ₂ CH ₂ SOMe	H	6-MeO-2-i-C ₃ F ₇	Amorphous
Q27	27-245	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-MeS-2-i-C ₃ F ₇	179
Q27	27-246	3-I	H		i-Pr	H 6-MeS-2-i-C ₃ F ₇	219
Q27	27-247	3-I	H		i-Pr	H 6-MeSO-2-i-C ₃ F ₇	Amorphous
Q27	27-248	3-I	H	C(Me) ₂ CH ₂ SMe	H	2-OCHF ₂	198
Q27	27-249	3-I	H	C(Me) ₂ CH ₂ SO ₂ Me	H	2-OCHF ₂	207

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-250	3-I	H	i-Pr	H	2-OCHF ₂	205
Q27	27-251	3-I	H	C (Me) ₂ CH ₂ SMe	H	2-SCHF ₂	174
Q27	27-252	3-I	H	i-Pr	H	2-SCHF ₂	226
Q27	27-253	3-I	H	i-Pr	H	2-SO ₂ CHF ₂	230
Q27	27-254	3-I	H	i-Pr	H	6-Me-2-OCHF ₂	252
Q27	27-255	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCHF ₂	124
Q27	27-256	3-I	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCHF ₂	185
Q27	27-257	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCHF ₂	102
Q27	27-258	3-I	H	i-Pr	H	6-Me-2-SCHF ₂	226
Q27	27-259	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-SCHF ₂	198
Q27	27-260	3-I	H	i-Pr	H	6-Me-2-OCH (CF ₃) ₂	266
Q27	27-261	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	223
Q27	27-262	3-I	H	i-Pr	H	6-Cl-2-OCH (CF ₃) ₂	216
Q27	27-263	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Cl-2-OCH (CF ₃) ₂	100
Q27	27-264	3-I	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	168
Q27	27-265	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	134
Q27	27-266	3-I	H	C (Me) ₂ CH ₂ SOMe	H	6-Cl-2-OCH (CF ₃) ₂	
Q27	27-267	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Cl-2-OCH (CF ₃) ₂	121
Q27	27-268	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-OMe-2-OCH (CF ₃) ₂	159
Q27	27-269	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-F-2-OCH (CF ₃) ₂	
Q27	27-270	3-I	H	i-Pr	H	OCH (CF ₃) ₂	240
Q27	27-271	3-I	H	t-Bu	H	OCH (CF ₃) ₂	
Q27	27-272	3-I	H	CH (Me) CH ₂ SMe	H	OCH (CF ₃) ₂	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-273	3-I	H	C (Me) ₂ CH ₂ SMe	H	OCH (CF ₃) ₂	237
Q27	27-274	3-I	H	i-Pr	H	2-Me-6-OCH (CF ₃) ₂	232
Q27	27-275	3-I	H	C (Me) ₂ CH ₂ SMe	H	2-Me-6-OCH (CF ₃) ₂	171
Q27	27-276	3-Cl	H	i-Pr	H	4-Me-2-OCH (CF ₃) ₂	226
Q27	27-277	3-I	H	i-Pr	H	4-Me-2-OCH (CF ₃) ₂	248
Q27	27-278	3-I	H	C (Me) ₂ CH ₂ SMe	H	4-Me-2-OCH (CF ₃) ₂	200
Q27	27-279	3-I	H	C (Me) ₂ CH ₂ SOMe	H	4-Me-2-OCH (CF ₃) ₂	118
Q27	27-280	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	4-Me-2-OCH (CF ₃) ₂	112
Q27	27-281	3-I	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-282	3-I	H	CH (Me) CH ₂ SEt	H	6-Me-2-OCH (CF ₃) ₂	256
Q27	27-283	H	H	i-Pr	H	6-Me-2-OCH (CF ₃) ₂	235
Q27	27-284	H	H	t-Bu	H	6-Me-2-OCH (CF ₃) ₂	255
Q27	27-285	H	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-286	H	H	CH (Me) CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-287	H	H	CH (Me) CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-288	H	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-289	H	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	108
Q27	27-290	H	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-291	3-F	H	i-Pr	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-292	3-F	H	t-Bu	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-293	3-F	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-294	3-F	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-295	3-F	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-296	3-F	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-297	3-Cl	H	i-Pr	H	6-Me-2-OCH (CF ₃) ₂	257
Q27	27-298	3-Cl	H	t-Bu	H	6-Me-2-OCH (CF ₃) ₂	255
Q27	27-299	3-Cl	Et	Et	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-300	3-Cl	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-301	3-Cl	H	CH (Me) CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-302	3-Cl	H	CH (Me) CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-303	3-Cl	H	CH (Me) CH ₂ SEt	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-304	3-Cl	H	CH (Me) CH ₂ SO ₂ Et	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-305	3-Cl	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	236
Q27	27-306	3-Cl	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	115
Q27	27-307	3-Cl	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	221
Q27	27-308	3-Br	H	i-Pr	H	6-Me-2-OCH (CF ₃) ₂	252
Q27	27-309	3-Br	H	t-Bu	H	6-Me-2-OCH (CF ₃) ₂	255
Q27	27-310	3-Br	Et	Et	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-311	3-Br	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-312	3-Br	H	CH (Me) CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-313	3-Br	H	CH (Me) CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-314	3-Br	H	CH (Me) CH ₂ SEt	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-315	3-Br	H	CH (Me) CH ₂ SO ₂ Et	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-316	3-Br	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	228
Q27	27-317	3-Br	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	115
Q27	27-318	3-Br	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	225

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-319	3-I	H		Me	H 6-Me-2-OCH(CF ₃) ₂	
Q27	27-320	3-I	H		Et	H 6-Me-2-OCH(CF ₃) ₂	
Q27	27-321	3-I	H		n-Pr	H 6-Me-2-OCH(CF ₃) ₂	
Q27	27-322	3-I	H		c-C ₃ H ₅	H 6-Me-2-OCH(CF ₃) ₂	
Q27	27-323	3-I	H		n-Bu	H 6-Me-2-OCH(CF ₃) ₂	261
Q27	27-324	3-I	H		s-Bu	H 6-Me-2-OCH(CF ₃) ₂	274
Q27	27-325	3-I	H		t-Bu	H 6-Me-2-OCH(CF ₃) ₂	241
Q27	27-326	3-I	H		i-Bu	H 6-Me-2-OCH(CF ₃) ₂	264
Q27	27-327	3-I	Et		Et	H 6-Me-2-OCH(CF ₃) ₂	165
Q27	27-328	3-I	Me		i-Pr	H 6-Me-2-OCH(CF ₃) ₂	
Q27	27-329	3-Cl-4-F	H		i-Pr	H 6-Me-2-OCH(CF ₃) ₂	
Q27	27-330	3-Cl-4-F	H	CH(Me)CH ₂ SMe	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-331	3-Cl-4-F	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-332	3,4-Cl ₂	H		i-Pr	H 6-Me-2-OCH(CF ₃) ₂	267
Q27	27-333	3,4-Cl ₂	H		t-Bu	H 6-Me-2-OCH(CF ₃) ₂	
Q27	27-334	3,4-Cl ₂	H	CH(Me)CH ₂ SMe	H	6-Me-2-OCH(CF ₃) ₂	210
Q27	27-335	3,4-Cl ₂	H	CH(Me)CH ₂ SOMe	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-336	3,4-Cl ₂	H	CH(Me)CH ₂ SO ₂ Me	H	6-Me-2-OCH(CF ₃) ₂	126
Q27	27-337	3,4-Cl ₂	H	CH(Me)CH ₂ SEt	H	6-Me-2-OCH(CF ₃) ₂	205
Q27	27-338	3,4-Cl ₂	H	CH(Me)CH ₂ SOEt	H	6-Me-2-OCH(CF ₃) ₂	119
Q27	27-339	3,4-Cl ₂	H	CH(Me)CH ₂ SO ₂ Et	H	6-Me-2-OCH(CF ₃) ₂	111
Q27	27-340	3,4-Cl ₂	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-341	3-Br-4-Cl	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-OCH(CF ₃) ₂	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-342	3,4-Br ₂	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-343	3-I-4-F	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-344	3-I-4-Cl	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-345	3-I-4-Br	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-346	3,4-I ₂	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-347	3-NO ₂	H		i-Pr	H	6-Me-2-OCH (CF ₃) ₂ 207
Q27	27-348	3-NO ₂	H		t-Bu	H	6-Me-2-OCH (CF ₃) ₂
Q27	27-349	3-NO ₂	Et		Et	H	6-Me-2-OCH (CF ₃) ₂
Q27	27-350	3-NO ₂	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-351	3-NO ₂	H	CH (Me) CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-352	3-NO ₂	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	244
Q27	27-353	3-NO ₂	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	230
Q27	27-354	3-CF ₃	H		i-Pr	H	6-Me-2-OCH (CF ₃) ₂ 211
Q27	27-355	3-CF ₃	H		t-Bu	H	6-Me-2-OCH (CF ₃) ₂ 246
Q27	27-356	3-CF ₃	Et		Et	H	6-Me-2-OCH (CF ₃) ₂
Q27	27-357	3-CF ₃	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-358	3-CF ₃	H	CH (Me) CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-359	3-CF ₃	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	226
Q27	27-360	3-CF ₃	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCH (CF ₃) ₂	112
Q27	27-361	3-CF ₃	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-362	3-OCF ₃	H		i-Pr	H	6-Me-2-OCH (CF ₃) ₂
Q27	27-363	3-OCF ₃	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	
Q27	27-364	3-OCF ₃	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH (CF ₃) ₂	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-365	3-SCF ₃	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-366	3-SCF ₃	H	CH(Me)CH ₂ SMe	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-367	3-SCF ₃	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-368	3-SOCF ₃	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-369	3-SO ₂ CF ₃	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-370	3-Me	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-371	3-Et	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-372	5-t-Bu	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	280
Q27	27-373	3-C≡CH	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-374	3-C≡CCF ₃	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-375	3-C≡C-t-Bu	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-376	3-C≡C-SiMe ₃	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-377	3-C≡C-Ph	H	i-Pr	H	6-Me-2-OCH(CF ₃) ₂	
Q27	27-378	3-I	H	i-Pr	H	6-Me-2-OCF ₂ CHF ₂	217
Q27	27-379	3-I	H	t-Bu	H	6-Me-2-OCF ₂ CHF ₂	
Q27	27-380	3-I	Et	Et	H	6-Me-2-OCF ₂ CHF ₂	
Q27	27-381	3-I	H	CH(Me)CH ₂ SMe	H	6-Me-2-OCF ₂ CHF ₂	
Q27	27-382	3-I	H	CH(Me)CH ₂ SOMe	H	6-Me-2-OCF ₂ CHF ₂	
Q27	27-383	3-I	H	CH(Me)CH ₂ SO ₂ Me	H	6-Me-2-OCF ₂ CHF ₂	
Q27	27-384	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-OCF ₂ CHF ₂	99
Q27	27-385	3-I	H	C(Me) ₂ CH ₂ SOMe	H	6-Me-2-OCF ₂ CHF ₂	
Q27	27-386	3-I	H	C(Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCF ₂ CHF ₂	
Q27	27-387	3-I	H	i-Pr	H	6-Cl-2-OCF ₂ CHF ₂	200

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-388	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Cl-2-OCF ₂ CHF ₂	142
Q27	27-389	3-I	H	C (Me) ₂ CH ₂ SOMe	H	6-Cl-2-OCF ₂ CHF ₂	
Q27	27-390	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Cl-2-OCF ₂ CHF ₂	
Q27	27-391	3-I	H	i-Pr	H	6-Me-2-OCF ₂ CHF ₂ CF ₃	205
Q27	27-392	3-I	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCF ₂ CHF ₂ CF ₃	
Q27	27-393	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCF ₂ CHF ₂ CF ₃	158
Q27	27-394	3-I	H	i-Pr	H	6-Me-2-OCF ₂ CHF ₂ CF ₃	
Q27	27-395	3-I	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCF ₂ CHF ₂ CF ₃	
Q27	27-396	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCF ₂ CHF ₂ CF ₃	126
Q27	27-397	3-I	H	i-Pr	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	194
Q27	27-398	3-I	H	t-Bu	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-399	3-I	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-400	3-I	H	CH (Me) CH ₂ SOMe	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-401	3-I	H	CH (Me) CH ₂ SO ₂ Me	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-402	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	91
Q27	27-403	3-I	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	81
Q27	27-404	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-OCF ₂ CHFOC ₃ F ₇ -n	157
Q27	27-405	3-I	H	i-Pr	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	205
Q27	27-406	3-I	H	t-Bu	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-407	3-I	H	CH (Me) CH ₂ SMe	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	106
Q27	27-408	3-I	H	CH (Me) CH ₂ SOMe	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-409	3-I	H	CH (Me) CH ₂ SO ₂ Me	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-410	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-411	3-I	H	C (Me) ₂ CH ₂ SOMe	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-412	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Cl-2-OCF ₂ CHFOC ₃ F ₇ -n	
Q27	27-413	3-I	H	i-Pr	H	6-Me-2-OCH ₂ C ₂ F ₅	259
Q27	27-414	3-I	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH ₂ C ₂ F ₅	208
Q27	27-415	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH ₂ C ₂ F ₅	
Q27	27-416	3-I	H	i-Pr	H	6-Me-2-OCH ₂ -n-C ₃ F ₇	
Q27	27-417	3-I	H	CH (Me) CH ₂ SMe	H	6-Me-2-OCH ₂ -n-C ₃ F ₇	
Q27	27-418	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-OCH ₂ -n-C ₃ F ₇	
Q27	27-419	3-I	H	i-Pr	H	6-Me-2-O- (2,4-Cl ₂ -Ph)	
Q27	27-420	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-O- (2,4-Cl ₂ -Ph)	
Q27	27-421	3-I	H	i-Pr	H	6-Me-2-O- (2-Cl-4-CF ₃ -Ph)	
Q27	27-422	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-O- (2-Cl-4-CF ₃ -Ph)	
Q27	27-423	3-I	H	i-Pr	H	6-Me-2-SCF ₃	
Q27	27-424	3-I	H	CH (Me) CH ₂ SMe	H	6-Me-2-SCF ₃	
Q27	27-425	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-SCF ₃	
Q27	27-426	3-I	H	C (Me) ₂ CH ₂ SOMe	H	6-Me-2-SCF ₃	
Q27	27-427	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-SCF ₃	
Q27	27-428	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-SOCF ₃	
Q27	27-429	3-I	H	C (Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-SO ₂ CF ₃	
Q27	27-430	3-I	H	i-Pr	H	6-Me-2-SC ₂ F ₅	
Q27	27-431	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-SC ₂ F ₅	
Q27	27-432	3-I	H	i-Pr	H	6-Me-2-S-n-C ₃ F ₇	
Q27	27-433	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-S-n-C ₃ F ₇	
Q27	27-436	3-I	H	C (Me) ₂ CH ₂ SMe	H	6-Me-2-S-CH (CF ₃) ₂	

Table 7 (Continued)


Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-434	3-I	H	i-Pr	H	6-Me-2-SCF ₂ CHF ₂	
Q27	27-435	3-I	H	CH(Me)CH ₂ SMe	H	6-Me-2-SCF ₂ CHF ₂ CF ₃	
Q27	27-437	3-I	H	i-Pr	H	6-Me-2-NHCOCF ₃	
Q27	27-438	3-I	H	i-Pr	H	6-Me-2-NHCOC ₂ F ₅	192
Q27	27-439	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-NHCOC ₂ F ₅	205
Q27	27-440	3-I	H	i-Pr	H	6-Me-2-NHCOC ₃ F _{7-n}	
Q27	27-441	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-NHCOC ₃ F _{7-n}	
Q27	27-442	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-NHCO- (2, 4-Cl ₂ -Ph)	
Q27	27-443	3-I	H	i-Pr	H	6-Me-2-NHCO- (4-CF ₃ -Ph)	
Q27	27-444	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-N(COC ₂ F ₅) ₂	
Q27	27-445	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-NHCH ₂ CF ₃	
Q27	27-446	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-NHCH ₂ CF ₃	
Q27	27-447	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-NHCH ₂ C ₂ F ₅	
Q27	27-448	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-Cl	173
Q27	27-449	3-CF ₃	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-i-C ₃ F ₇	214
Q27	27-450	H	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-i-C ₃ F ₇	155
Q27	27-451	3-F	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-i-C ₃ F ₇	202
Q27	27-452	3-F	H	C(Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-i-C ₃ F ₇	197
Q27	27-453	3-Br	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-i-C ₃ F ₇	206
Q27	27-454	3-Br	H	C(Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-i-C ₃ F ₇	225
Q27	27-455	3, 4-Cl ₂	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-i-C ₃ F ₇	259
Q27	27-456	3-CF ₃	H	i-Pr	H	6-Me-2-i-C ₃ F ₇	221

Table 7 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q27	27-457	H	H	i-Pr	H	6-Cl-2-i-C ₃ F ₇	200
Q27	27-458	H	H	C(Me) ₂ CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	110
Q27	27-459	3-Cl	H	i-Pr	H	6-Cl-2-i-C ₃ F ₇	181
Q27	27-460	3-I	H	t-Bu	H	6-Cl-2-i-C ₃ F ₇	191
Q27	27-461	3-F	H	C(Me) ₂ CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	182
Q27	27-462	3-Cl	H	C(Me) ₂ CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	161
Q27	27-463	3,4-Cl ₂	H	C(Me) ₂ CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	153
Q27	27-464	3-CF ₃	H	C(Me) ₂ CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	165
Q27	27-465	3-NO ₂	H	C(Me) ₂ CH ₂ SMe	H	6-Cl-2-i-C ₃ F ₇	235
Q27	27-466	3-I	H	i-Pr	H	6-Me-2-O(4-CF ₃ -Ph)	238
Q27	27-467	3-I	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-O(4-CF ₃ -Ph)	111
Q27	27-468	3-I	H	C(Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-O(4-CF ₃ -Ph)	106
Q27	27-469	3-I	H	C(Me) ₂ CH ₂ SO ₂ Me	H	6-Me-2-O(4-CF ₃ -Ph)	97
Q27	27-470	3-CF ₃	H	C(Me) ₂ CH ₂ SMe	H	6-Me-2-OCF ₂ CHFOCF ₃	
							Amorphous
Q27	27-471	3-I	H	i-Pr	H	6-Me-2-OCF=CF ₂ CF ₃	165
Q27	27-472	3-I	H	i-Pr	H	6-Me-2-OCF ₂ CHFOCF ₃	185

In Table 7, some compounds are amorphous or pasty. ^1H -NMR data of such compounds are shown below.

No	^1H -NMR [δ (ppm/ CDCl_3)]
27-153	1.59(s. 3H), 1.64(s. 3H), 2.26(s. 3H), 2.62(s. 3H), 2.88(d. 1H), 3.02(d. 1H), 6.83(br. 1H), 7.23(t. 1H), 7.58(dd. 1H), 7.78(d. 1H), 8.00(dd. 1H), 8.58(br. 1H), 8.81(d. 1H).
27-235	1.64(s. 3H), 1.66(s. 3H), 2.40(s. 3H), 2.88(d. 1H), 3.24(d. 1H), 6.72(br. 1H), 7.24(t. 1H), 7.70(dd. 1H), 7.74(d. 1H), 8.03(dd. 1H), 8.85(br. 1H), 9.12(d. 1H).
27-236	1.71(s. 6H), 2.71(s. 3H), 3.63(s. 2H), 6.25(br. 1H), 7.25(t. 1H), 7.70(dd. 1H), 7.75(dd. 1H), 8.05(dd. 1H), 8.81(br. 1H), 9.11(d. 1H).
27-238	1.68(s. 3H), 1.72(s. 3H), 2.49(s. 3H), 2.99(d. 1H), 3.21(d. 1H), 6.76(br. 1H), 7.21(t. 1H), 7.50(dd. 1H), 7.66(dd. 1H), 7.84(dd. 1H), 8.37(dd. 1H), 8.68(d. 1H), 9.75(br. 1H).
27-239	1.80(s. 6H), 2.87(s. 3H), 3.73(s. 2H), 6.23(br. 1H), 7.23(t. 1H), 7.43(dd. 1H), 7.65(dd. 1H), 7.82(dd. 1H), 8.35(dd. 1H), 8.64(d. 1H), 9.88(br. 1H).
27-244	1.60(s. 3H), 1.63(s. 3H), 2.41(s. 3H), 2.84(d. 1H), 3.31(d. 1H), 4.02(s. 1H), 6.59(br. 1H), 7.21(t. 1H), 7.34(dd. 1H), 7.69(dd. 1H), 7.99(dd. 1H), 8.65(br. 1H), 8.88(d. 1H).
27-247	1.31(dd. 6H), 3.50(s. 3H), 4.33(m. 1H), 5.60(d. 1H), 7.19(t. 1H), 7.68(d. 1H), 7.74(dd. 1H), 8.00(d. 1H), 9.26(d. 1H), 11.8(br. 1H).
27-470	1.42(s. 6H), 1.96(s. 3H), 2.53(s. 3H), 2.81(s. 2H), 6.17(s. 1H), 6.62(dt. 1H), 6.90(d. 1H), 7.66(t. 1H), 7.85(d. 1H), 8.03(d. 1H), 8.63(d. 1H), 8.71(s. 1H).

Table 8 ($Z^1 = Z^2 = O$)


Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q29	29- 1	3-Cl	H	i-Pr	H	H	
Q29	29- 2	3-Cl	H	i-Pr	H	4-Me-6-C ₂ F ₅	
Q29	29- 3	3-Br	H	i-Pr	H	4-Me-6-n-C ₃ F ₇	
Q29	29- 4	3-NO ₂	H	i-Pr	H	4-Me-6-i-C ₃ F ₇	
Q29	29- 5	3-I	H	i-Pr	H	H	
Q29	29- 6	3-I	H	i-Pr	H	6-Cl	136
Q29	29- 7	3-I	H	i-Pr	H	6-C ₂ F ₅	
Q29	29- 8	3-I	H	i-Pr	H	6-n-C ₃ F ₇	
Q29	29- 9	3-I	H	i-Pr	H	6-i-C ₃ F ₇	
Q29	29-10	3-I	H	i-Pr	H	4-Me-6-C ₂ F ₅	
Q29	29-11	3-I	H	i-Pr	H	4-Me-6-n-C ₃ F ₇	
Q29	29-12	3-I	H	i-Pr	H	4-Me-6-i-C ₃ F ₇	
Q29	29-13	3-I	H	i-Pr	H	4-Me-5-C ₂ F ₅	
Q29	29-14	3-I	H	i-Pr	H	4-Me-5-n-C ₃ F ₇	
Q29	29-15	3-I	H	i-Pr	H	4-Me-5-i-C ₃ F ₇	

Table 8 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q29	29-16	3-I	H	t-Bu	H	6-i-C ₃ F ₇	
Q29	29-17	3-I	H	t-Bu	H	6-C ₂ F ₅	
Q29	29-18	3-I	H	t-Bu	H	6-n-C ₃ F ₇	
Q29	29-19	3-I	H	t-Bu	H	4-Me-6-i-C ₃ F ₇	
Q29	29-20	3-I	H	CH(CH ₃)CH ₂ SCH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-21	3-I	H	CH(CH ₃)CH ₂ SOCH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-22	3-I	H	CH(CH ₃)CH ₂ SO ₂ CH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-23	3-I	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-24	3-I	H	C(CH ₃) ₂ CH ₂ SOCH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-25	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-26	3-I	H	CH(CH ₃)CH ₂ NHAc	H	4-Me-6-i-C ₃ F ₇	
Q29	29-27	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	4-Me-6-i-C ₃ F ₇	
Q29	29-28	3-I	H	CH(CH ₃)CH ₂ CH ₂ OCH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-29	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	4-Me-6-i-C ₃ F ₇	
Q29	29-30	3-I	Et	Et	H	H	
Q29	29-31	3-I	Et	Et	H	4-Me-6-C ₂ F ₅	
Q29	29-32	3-I	Et	Et	H	4-Me-6-n-C ₃ F ₇	
Q29	29-33	3-I	Et	Et	H	4-Me-6-i-C ₃ F ₇	
Q29	29-34	3-I	Et	Et	H	6-Cl	
Q29	29-35	3-I	Et	Et	H	6-Br	
Q29	29-36	3-I	Et	Et	H	6-n-C ₃ F ₇	
Q29	29-37	3-CF ₃	H	i-Pr	H	4-Me-6-C ₂ F ₅	
Q29	29-38	3-Ph	H	i-Pr	H	4-Me-6-n-C ₃ F ₇	

Table 8 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q29	29-39	3-SOCF ₃	H	i-Pr	H	4-Me-6-i-C ₃ F ₇	
Q29	29-40	3-C ₂ F ₅	H	i-Pr	H	4-Me-6-C ₂ F ₅	
Q29	29-41	3-I-4-Cl	H	i-Pr	H	4-Me-6-n-C ₃ F ₇	
Q29	29-42	3-I-4-CF ₃	H	i-Pr	H	4-Me-6-i-C ₃ F ₇	
Q29	29-43	3-CF ₃ -4-Cl	H	i-Pr	H	4-Me-6-C ₂ F ₅	
Q29	29-44	3-OCF ₂ O-4	H	i-Pr	H	4-Me-6-n-C ₃ F ₇	
Q29	29-45	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	4-Me-6-i-C ₃ F ₇	



Q32


Table 9 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q32	32- 1	3-Cl	H i-Pr		H H		
Q32	32- 2	3-Cl	H i-Pr		H	4-Me-2-C ₂ F ₅	
Q32	32- 3	3-Br	H i-Pr		H	4-Me-2-n-C ₃ F ₇	
Q32	32- 4	3-NO ₂	H i-Pr		H	4-Me-2-i-C ₃ F ₇	
Q32	32- 5	3-I	H i-Pr		H	2-C ₂ F ₅	
Q32	32- 6	3-I	H i-Pr		H	2-n-C ₃ F ₇	
Q32	32- 7	3-I	H i-Pr		H	2-i-C ₃ F ₇	
Q32	32- 8	3-I	H i-Pr		H	4-Me-2-C ₂ F ₅	
Q32	32- 9	3-I	H i-Pr		H	4-Me-2-n-C ₃ F ₇	
Q32	32-10	3-I	H i-Pr		H	4-Me-2-i-C ₃ F ₇	
Q32	32-11	3-I	H i-Pr		H	4,6-Cl ₂	257
Q32	32-12	3-I	H t-Bu		H	2-i-C ₃ F ₇	
Q32	32-13	3-I	H t-Bu		H	2-C ₂ F ₅	
Q32	32-14	3-I	H t-Bu		H	2-n-C ₃ F ₇	
Q32	32-15	3-I	H t-Bu		H	4-Me-2-i-C ₃ F ₇	
Q32	32-16	3-I	H CH(CH ₃)CH ₂ SCH ₃		H	4-Me-2-i-C ₃ F ₇	
Q32	32-17	3-I	H CH(CH ₃)CH ₂ SOCH ₃		H	4-Me-2-i-C ₃ F ₇	
Q32	32-18	3-I	H CH(CH ₃)CH ₂ SO ₂ CH ₃		H	4-Me-2-i-C ₃ F ₇	
Q32	32-19	3-I	H C(CH ₃) ₂ CH ₂ SCH ₃		H	4-Me-2-i-C ₃ F ₇	202
Q32	32-20	3-I	H C(CH ₃) ₂ CH ₂ SOCH ₃		H	4-Me-2-i-C ₃ F ₇	
Q32	32-21	3-I	H C(CH ₃) ₂ CH ₂ SO ₂ CH ₃		H	4-Me-2-i-C ₃ F ₇	
Q32	32-22	3-I	H CH(CH ₃)CH ₂ SEt		H	4-Me-2-i-C ₃ F ₇	
Q32	32-23	3-I	H C(CH ₃) ₂ CH ₂ SEt		H	4-Me-2-i-C ₃ F ₇	

[illegible]

Amorphous

No	$^1\text{H-NMR}$ [δ (ppm/ CDCl_3)]
32-42	1.04 (s. 3H), 1.31 (t. 3H), 3.10 (m. 3H), 3.42 (m. 1H), 3.80 (m. 1H), 4.96-4.74 (m. 4H), 7.22 (t. 1H), 7.87 (d. 1H), 8.04 (dd. 1H), 8.39 (s. 1H).

Table 10 ($Z^1 = Z^2 = O$)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C), nD (°C)
Q33	33- 1	H	H i-Pr		H	4, 6- (OMe) ₂	61
Q33	33- 2	3-Cl	H i-Pr		H	H	
Q33	33- 3	3-Cl	H i-Pr		H	5-C ₂ F ₅	
Q33	33- 4	3-Br	H i-Pr		H	5-n-C ₃ F ₇	
Q33	33- 5	3-NO ₂	H i-Pr		H	5-i-C ₃ F ₇	
Q33	33- 6	3-I	H i-Pr		H	5-C ₂ F ₅	
Q33	33- 7	3-I	H i-Pr		H	5-n-C ₃ F ₇	
Q33	33- 8	3-I	H i-Pr		H	5-i-C ₃ F ₇	
Q33	33- 9	3-I	H i-Pr		H	4, 6-OMe ₂	nD 1.5672 (20.9)
Q33	33-10	3-I	H i-Pr		H	4, 6-OMe ₂ -5-i-C ₃ F ₇	nD 1.5045 (21.9)
Q33	33-11	3-I	H t-Bu		H	5-i-C ₃ F ₇	
Q33	33-12	3-I	H t-Bu		H	5-C ₂ F ₅	
Q33	33-13	3-I	H t-Bu		H	5-n-C ₃ F ₇	
Q33	33-14	3-I	H CH (CH ₃) CH ₂ SCH ₃		H	5-i-C ₃ F ₇	
Q33	33-15	3-I	H CH (CH ₃) CH ₂ SOCH ₃		H	5-i-C ₃ F ₇	

Table 10 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C), nD (°C)
Q33	33-16	3-I	H	CH(CH ₃)CH ₂ SO ₂ CH ₃	H	5-i-C ₃ F ₇	
Q33	33-17	3-I	H	C(CH ₃) ₂ CH ₂ SCH ₃	H	5-i-C ₃ F ₇	
Q33	33-18	3-I	H	C(CH ₃) ₂ CH ₂ SOCH ₃	H	5-i-C ₃ F ₇	
Q33	33-19	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	5-i-C ₃ F ₇	
Q33	33-20	3-I	H	CH(CH ₃)CH ₂ SEt	H	5-i-C ₃ F ₇	
Q33	33-21	3-I	H	C(CH ₃) ₂ CH ₂ SEt	H	5-i-C ₃ F ₇	
Q33	33-22	3-I	H	CH(CH ₃)CH ₂ CH ₂ SCH ₃	H	5-i-C ₃ F ₇	
Q33	33-23	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ SCH ₃	H	5-i-C ₃ F ₇	
Q33	33-24	3-I	Et	Et	H	5-C ₂ F ₅	
Q33	33-25	3-I	Et	Et	H	5-n-C ₃ F ₇	
Q33	33-26	3-I	Et	Et	H	5-i-C ₃ F ₇	
Q33	33-27	3-I	Et	Et	H	5-Cl	
Q33	33-28	3-I	Et	Et	H	5-Br	
Q33	33-29	3-I	Et	Et	H	5-n-C ₃ F ₇	




Table 11 ($Z^1 = Z^2 = O$)



Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q34	34- 1	3-Cl	H i-Pr		H H		
Q34	34- 2	3-Cl	H i-Pr		H 3-Me-5-C ₂ F ₅		
Q34	34- 3	3-Br	H i-Pr		H 3-Me-5-n-C ₃ F ₇		
Q34	34- 4	3-NO ₂	H i-Pr		H 3-Me-5-i-C ₃ F ₇		
Q34	34- 5	3-I	H i-Pr		H H		185
Q34	34- 6	3-I	H i-Pr		H 5-I		198
Q34	34- 7	3-I	H i-Pr		H 5-C ₂ F ₅		
Q34	34- 8	3-I	H i-Pr		H 5-n-C ₃ F ₇		
Q34	34- 9	3-I	H i-Pr		H 5-i-C ₃ F ₇		
Q34	34-10	3-I	H i-Pr		H 3-Me-5-C ₂ F ₅		
Q34	34-11	3-I	H i-Pr		H 3-Me-5-n-C ₃ F ₇		
Q34	34-12	3-I	H i-Pr		H 3-Me-5-i-C ₃ F ₇		
Q34	34-13	3-I	H i-Pr		H 6-Me-5-C ₂ F ₅		
Q34	34-14	3-I	H i-Pr		H 6-Me-5-n-C ₃ F ₇		
Q34	34-15	3-I	H i-Pr		H 6-Me-5-i-C ₃ F ₇		
Q34	34-16	3-I	H t-Bu		H 5-i-C ₃ F ₇		
Q34	34-17	3-I	H t-Bu		H 5-C ₂ F ₅		
Q34	34-18	3-I	H t-Bu		H 5-n-C ₃ F ₇		
Q34	34-19	3-I	H t-Bu		H 3-Me-5-i-C ₃ F ₇		
Q34	34-20	3-I	H CH(CH ₃)CH ₂ SCH ₃		H 3-Me-5-i-C ₃ F ₇		
Q34	34-21	3-I	H CH(CH ₃)CH ₂ SOCH ₃		H 3-Me-5-i-C ₃ F ₇		
Q34	34-22	3-I	H CH(CH ₃)CH ₂ SO ₂ CH ₃		H 3-Me-5-i-C ₃ F ₇		
Q34	34-23	3-I	H C(CH ₃) ₂ CH ₂ SCH ₃		H 3-Me-5-i-C ₃ F ₇		

Table 11 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q34	34-24	3-I	H	C(CH ₃) ₂ CH ₂ SOCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q34	34-25	3-I	H	C(CH ₃) ₂ CH ₂ SO ₂ CH ₃	H	3-Me-5-i-C ₃ F ₇	
Q34	34-26	3-I	H	CH(CH ₃)CH ₂ NHAc	H	3-Me-5-i-C ₃ F ₇	
Q34	34-27	3-I	H	C(CH ₃) ₂ CH ₂ NHAc	H	3-Me-5-i-C ₃ F ₇	
Q34	34-28	3-I	H	CH(CH ₃)CH ₂ CH ₂ OCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q34	34-29	3-I	H	C(CH ₃) ₂ CH ₂ CH ₂ OCH ₃	H	3-Me-5-i-C ₃ F ₇	
Q34	34-30	3-I	Et	Et	H	H	144
Q34	34-31	3-I	Et	Et	H	3-Me-5-C ₂ F ₅	
Q34	34-32	3-I	Et	Et	H	3-Me-5-n-C ₃ F ₇	
Q34	34-33	3-I	Et	Et	H	3-Me-5-i-C ₃ F ₇	
Q34	34-34	3-I	Et	Et	H	5-Cl	
Q34	34-35	3-I	Et	Et	H	5-Br	
Q34	34-36	3-I	Et	Et	H	5-n-C ₃ F ₇	
Q34	34-37	3-CF ₃	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q34	34-38	3-Ph	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q34	34-39	3-SOCF ₃	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	
Q34	34-40	3-C ₂ F ₅	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q34	34-41	3-I-4-Cl	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q34	34-42	3-I-4-CF ₃	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	
Q34	34-43	3-CF ₃ -4-Cl	H	i-Pr	H	3-Me-5-C ₂ F ₅	
Q34	34-44	3-OCF ₂ O-4	H	i-Pr	H	3-Me-5-n-C ₃ F ₇	
Q34	34-45	3-OCF ₂ CF ₂ O-4	H	i-Pr	H	3-Me-5-i-C ₃ F ₇	
Q34	34-46	3-I	H	Et	H	5-i-C ₃ F ₇	175

Table 11 (Continued)

Q	No.	Xn	R ¹	R ²	R ³	Ym	mp (°C)
Q3	3- 1	3-I	H	i-Pr	H	H	
Q7	7- 1	3-I	H	i-Pr	H	H	
Q11	11- 1	3-I	H	i-Pr	H	H	
Q14	14- 1	3-I	H	i-Pr	H	H	
Q15	15- 1	3-I	H	i-Pr	H	H	185
Q18	18- 1	3-I	H	i-Pr	H	H	
Q20	20- 1	3-I	H	i-Pr	H	H	

Table 12 (Z¹ = Z² = O)

Q	No.	Xn	R ¹	R ²	R ³	W	Ym	Property, Mp (°C)
Q16	16- 1	3-Cl	H	i-Pr	H	N-i-Pr	SMe	Paste
Q16	16- 2	3-Cl	H	i-Pr	H	N-n-Pr	SMe	Paste
Q44	43- 1	H	H	i-Pr	H	S	6-Cl	47

EXAMPLES

Next, typical examples of the present invention are shown below. The invention is by no means limited by these examples.

5 Production Example 1

(1-1) Production of N-(4-methyl-3-trifluoromethyl-isoxazol-5-yl)-3-iodophthalimide

In 20 ml of acetic acid, 0.6 g of 3-iodophthalic anhydride and 0.44 g of 5-amino-4-methyl-
10 3-trifluoromethyl-isoxazole were dissolved and reacted for 9 hours with heating under reflux. After completion of the reaction, the solvent was distilled off under reduced pressure, and the residue was dissolved in ethyl acetate, washed with dilute hydrochloric acid,
15 saturated aqueous solution of sodium bicarbonate and saturated aqueous solution of sodium chloride, and then dried on sodium sulfate. After distilling off the solvent under reduced pressure, the residue was purified by silica gel column chromatography using a
20 3/1 mixture of hexane and ethyl acetate as an eluent to obtain 0.71 g of the objective product.

Property: m.p. 105°C; Yield: 69%

(1-2) Production of N¹-(4-methyl-3-trifluoromethyl-isoxazol-5-yl)-N²-isopropyl-3-iodophthalamide (Compound
25 No. Q6-8)

In 50 ml of dioxane was dissolved 1.06 g of N-(4-methyl-3-trifluoromethylisoxazol-5-yl)-3-iodophthaimide. Then, 0.4 g of isopropylamine was

added to the solution obtained above and stirred at room temperature for 3 hours. After completion of the reaction, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel
5 column chromatography using a 2/1 mixture of hexane and ethyl acetate as an eluent to obtain 0.32 g of the objective product.

Property: m.p. 103°C; Yield: 26%

Production Example 2

10 (2-1) Production of N¹-[2-(1,1-dimethylethyl)-1,3,4-thiadiazol-4-yl]-N²,N²-diethyl-3-iodophthalamide (Compound No. Q19-34)

In 20 ml of tetrahydrofuran were dissolved 0.5 g of N,N-diethyl-3-iodophthalic acid 2-amide and
15 0.27 g of 5-amino-2-(1,1-dimethylethyl)-1,3,4-thiadiazole. After adding 0.28 g of diethylphosphoryl cyanide and 0.18 g of triethylamine, a reaction was carried out for 7 hours with heating under reflux. After completion of the reaction, ethyl acetate was
20 added to the reaction mixture, and the resulting ethyl acetate solution was washed with dilute hydrochloric acid, saturated aqueous solution of sodium bicarbonate and saturated aqueous solution of sodium chloride, and dried on sodium sulfate. After distilling off the
25 solvent, the residue was purified by silica gel column chromatography using 3/1 mixture of hexane and ethyl acetate to obtain 0.11 g of the objective product.

Property: m.p. 59°C; Yield: 16%

Production Example 3

(3-1) Production of N¹-(4-methyl-2-heptafluoroisopropyl-pyridin-5-yl)-N²-(1-methyl-2-methylthioethyl)-3-

5 iodophthalamide (Compound No. Q27-144)

In 4 ml of acetonitrile were dissolved 0.37 g of N-(1-methyl-2-methylthioethyl)-3-iodophthalic acid isoimide and 0.28 g of 5-amino-4-methyl-2-heptafluoro-isopropyl-pyridine. After adding a catalytic quantity
10 of trifluoroacetic acid, the resulting mixture was stirred at room temperature for 30 minutes. The resulting crystal was collected by filtration, and there was obtained 0.28 g of the objective product.

Property: m.p. 225°C; Yield: 44%

15 Production Example 4: Production of 2-amino-3-methyl-6-pentafluoroethylpyridine (Compound No. IV-1)

To 20 ml of dimethyl sulfoxide were added 2.34 g (0.01 mol) of 2-amino-5-iodo-3-methylpyridine, 2.5 g of powdered metallic copper and 3.7 g (0.015 mol)
20 of iodopentafluorethane. The mixture was kept at 110°C and vigorously stirred for 6 hours. After cooling the reaction mixture to room temperature, the mixture was poured into 500 ml of ice water and thoroughly stirred. The insoluble matter was filtered off, and the objec-
25 tive product was extracted from the filtrate with 300 ml of ethyl acetate. The extract solution was washed

with water, dried on anhydrous sodium sulfate, and concentrated under reduced pressure. Purification of the residue by column chromatography using 3/7 mixture of ethyl acetate and hexane as an eluent gave 1.1 g of the objective product (yield 20%).

$^1\text{H-NMR}$ [δ (CDCl_3)]: 2.17 (s, 3H), 4.82 (br, 2H), 7.42 (d, 1H), 8.16 (s, 1H),

Production Example 5: Production of 3-amino-2-methoxy-6-(heptafluoropropan-2-yl)-pyridine (Compound No. IV-15)

To 20 ml of methyl t-butyl ether were added 3.2 g (0.026 mol) of 3-amino-2-methoxypyridine, 0.6 g of triethylbenzylammonium chloride, 2.0 g of sodium carbonate and 10.0 g (0.031 mol) of 2-iodoheptafluoropropane. While stirring the mixture at 30°C, a solution of 2.8 of sodium hydrosulfite in 10 ml water was dropwise added thereto. After the dropping, the resulting mixture was reacted at room temperature for 20 hours, after which the organic layer was separated, washed with water and dried on anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. Purification of the residue by column chromatography using 3/7 mixture of ethyl acetate and hexane as an eluent gave 2.0 g of the objective product (yield 26%).

$^1\text{H-NMR}$ [δ (CDCl_3)]: 3.96 (s, 3H), 4.03 (br, 2H), 6.91 (d, 1H), 7.10 (dd, 1H)

Production Example 6: Production of 3-amino-6-(1,1,1,3,3,3-hexafluoroisopropoxy)pyridine (Compound No. IV-27)

Sodium hydride (2.6 g) was portionwise added to a solution of 11.3 g of 1,1,1,3,3,3-hexafluoro-2-propanol in 50 ml tetrahydrofuran at a temperature not exceeding 5°C, and stirred at the same temperature as above for 30 minutes. Then, 4.7 g of 2-chloro-5-nitropyridine was added and stirred at room temperature for 12 hours. The reaction mixture was poured into 200 ml of ice water and extracted with 300 ml of ethyl acetate. The extract solution was washed with water, dried on anhydrous sodium sulfate and concentrated. Purification of the residue by column chromatography using 1/10 mixture of ethyl acetate and hexane as an eluent gave 6.2 g of 2-(1,1,1,3,3,3-hexafluoroisopropoxy)-5-nitrobenzene (yield 64%).

To 20 ml of acetic acid were added 4.4 g of the 2-(1,1,1,3,3,3-hexafluoroisopropoxy)-5-nitrobenzene obtained above and 4.2 g of electrolytic iron. The resulting mixture was stirred at 60-65°C for 30 minutes. The reaction mixture was cooled and neutralized with 6N aqueous solution of sodium hydroxide, the insoluble matter was filtered off, and the filtrate was extracted with 300 ml of t-butyl methyl ether. The extract solution was washed with aqueous solution of sodium chloride, dried on anhydrous sodium sulfate and concentrated. Purification of the

residue by column chromatography using 2/3 mixture of ethyl acetate and hexane as an eluent gave 3.6 g of 3-amino-6-(1,1,1,3,3,3-hexafluoroisopropoxy)pyridine (yield 92%).

- 5 The agrohorticultural insecticides containing the phthalamide derivative of formula (I) of the present invention as an active ingredient are suitable for controlling various insect pests such as agrohorticultural insect pests, stored grain insect pests,
- 10 sanitary insect pests, nematodes, etc., which are injurious to paddy rice, fruit trees, vegetables, other crops, flowers, ornamental plants, etc. They have a marked insecticidal effect, for example, on LEPIDOPTERA including summer fruit tortrix (Adoxophyes orana
- 15 fasciata), smaller tea tortrix (Adoxophyes sp.), Manchurian fruit moth (Grapholita inopinata), oriental fruit moth (Grapholita molesta), soybean pod border (Leguminovora glycinivorella), mulberry leafroller (Olethreutes mori), tea leafroller (Caloptilia
- 20 thevivora), Caloptilia sp. (Caloptilia zachrysa), apple leafminer (Phyllonorycter ringoniella), pear barkminer (Splerrina astaurota), common white (Piers rapae crucivora), tobacco budworm (Heliothis sp.), codling moth (Laspey resia pomonella), diamondback moth
- 25 (Plutella xylostella), apple fruit moth (Argyresthia conjugella), peach fruit moth (Carposina niponensis), rice stem borer (Chilo suppressalis), rice leafroller (Cnaphalocrocis medinalis), tobacco moth (Ephestia

- elutella), mulberry pyralid (Glyphodes pyloalis), yellow rice borer (Scirpophaga incertulas), rice skipper (Parnara guttata), rice armyworm (Pseudaletia separata), pink borer (Sesamia inferens), common
- 5 cutworm (Spodoptera litura), beet armyworm (Spodoptera exigua), etc.; HEMIPTERA including aster leafhopper (Macrosteles fascifrons), green rice leafhopper (Nephotettix cincticeps), brown rice planthopper (Nilaparvata lugens), whitebacked rice planthopper
- 10 (Sogatella furcifera), citrus psylla (Diaphorina citri), grape whitefly (Aleurolibus taonabae), sweetpotato whitefly (Bemisia tabaci), greenhouse whitefly (Trialeurodes vaporariorum), turnup aphid (Lipaphis erysimi), green peach aphid (Myzus persicae),
- 15 Indian wax scale (Ceroplastes ceriferus), cottony citrus scale (Pulvinaria aurantii), camphor scale (Pseudaonidia duplex), san Jose scale (Comstockaspis perniciosa), arrowhead scale (Unapsis yanonensis), etc.; TYLENCHIDA including soybean beetle (Anomala
- 20 rufocuprea), Japanese beetle (Popillia japonica), tobacco beetle (Lasionderma serricorne), powderpost beetle (Lyctus brunneus), twenty-eight-spotted ladybird (Epilachna vigintiotopunctata), azuki bean weevil (Callosobruchus chinensis), vegetable weevil
- 25 (Listroderes costirostris), maize weevil (Sitophilus zeamais), boll weevil (Anthonomus grandis grandis), rice water weevil (Lissorhoptrus oryzophilus), cucurbit leaf beetle (Aulacophora femoralis), rice leaf beetle

(Oulema oryzae), striped flea beetle (Phyllotreta striolata), pine shoot beetle (Tomicus piniperda), Colorado potato beetle (Leptinotarsa decemlineata), Mexican bean beetle (Epilachna varivestis), corn
 5 rootworm (Diabrotica sp.), etc.; DIPTERA including (Dacus(Zeugodacus) cucurbitae), oriental fruit fly (Dacus(Bactrocera) dorsalis), rice leafminer (Agnomyza oryzae), onion maggot (Delia antiqua), seedcorn maggot (Delia platura), soybean pod gall midge (Asphondylia
 10 sp.), muscid fly (Musca domestica), house mosquito (Culex pipiens pipiens), etc.; and TYLENCHIDA including root-lesion nematode (Pratylenchus sp.), coffee root-lesion nematode (Pratylenchus coffeae), potato cyst nematode (Globodera rostochiensis), root-knot nematode
 15 (Meloidogyne sp.), citrus nematode (Tylenchulus semipenetrans), Aphelenchus sp. (Aphelenchus avenae), chrysanthemum foliar (Aphelenchoides ritzemabosi), etc.

The agrohorticultural agent and particularly the agrohorticultural insecticide containing the
 20 phthalamide derivative represented by formula (I) of the present invention has a marked controlling effect on the above-exemplified insect pests, sanitary pests and/or nematodes, which are injurious to paddy field crops, upland crops, fruit trees, vegetables and other
 25 crops, flowers and ornament plants, and the like. Therefore, the desired effect of the agrohorticultural insecticide of the present invention can be exhibited by applying the insecticide to the paddy field water,

stalks and leaves or soil of paddy field, upland field, fruit trees, vegetables, other crops or flowers and ornament plants at a season at which the insect pests, sanitary pests or nematodes are expected to appear, before their appearance or at the time when their appearance is confirmed.

In general, the agrohorticultural agent of the present invention is used after being prepared into conveniently usable forms according to ordinary manner for preparation of agrochemicals.

That is, the phthalamide derivative of formula (I) and an appropriate carrier are blended optionally together with an adjuvant in a proper proportion and prepared into a suitable preparation form such as suspension, emulsifiable concentrate, soluble concentrate, wettable powder, granules, dust or tablets through dissolution, separation, suspension, mixing, impregnation, adsorption or sticking.

The inert carrier used in the present invention may be either solid or liquid. As the solid carrier, soybean flour, cereal flour, wood flour, bark flour, saw dust, powdered tobacco stalks, powdered walnut shells, bran, powdered cellulose, extraction residues of vegetables, powdered synthetic polymers or resins, clay (e.g. kaolin, bentonite and acid clay), talc (e.g. talc and pyrophyllite), silica materials (e.g. diatomaceous earth, siliceous sand, mica, white carbon, i.e. synthetic high-dispersion silicic acid,

also called finely divided hydrated silica or hydrated silicic acid, some of the commercially available products contain calcium silicate as the major component), activated carbon, powdered sulfur, pumice, 5 calcined diatomaceous earth, ground brick, fly ash, sand, calcium carbonate, calcium phosphate and other inorganic or mineral powders, chemical fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, ammonium chloride and the like, and 10 compost. These carriers may be used either alone or as a mixture of two or more carriers.

The liquid carrier is that which itself has a solubility or which is without such solubility but is capable of dispersing an active ingredient with the aid 15 of an adjuvant. The following are typical examples of the liquid carrier and can be used alone or as a mixture thereof. Water; alcohols such as methanol, ethanol, isopropanol, butanol and ethylene glycol; ketones such as acetone, methyl ethyl ketone, methyl 20 isobutyl ketone, diisobutyl ketone and cyclohexanone; ethers such as ethyl ether, dioxane, cellosolve, dipropyl ether and tetrahydrofuran; aliphatic hydrocarbons such as kerosene and mineral oil; aromatic hydrocarbons such as benzene, toluene, xylene, solvent 25 naphtha and alkylnaphthalene; halogenated hydrocarbons such as dichlorethane, chloroform, carbon tetrachloride and chlorobenzene; esters such as ethyl acetate, diisopropyl phthalate, dibutyl phthalate and dioctyl

phthalate; amides such as dimethylformamide, diethylformamide and dimethylacetamide; nitriles such as acetonitrile; and dimethyl sulfoxide.

The following are typical examples of the
5 adjuvant, which are used depending upon purposes and used alone or in combination of two or more adjuvants in some cases, or need not to be used at all.

To emulsify, disperse, dissolve and/or wet an active ingredient, a surfactant is used. As the
10 surfactant, there can be exemplified polyoxyethylene alkyl ethers, polyoxyethylene alkylaryl ethers, polyoxyethylene higher fatty acid esters, polyoxyethylene resinsates, polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monooleate,
15 alkylarylsulfonates, naphthalene-sulfonic acid condensation products, ligninsulfonates and higher alcohol sulfate esters.

Further, to stabilize the dispersion of an active ingredient, tackify it and/or bind it, there may
20 be used adjuvants such as casein, gelatin, starch, methyl cellulose, carboxymethyl cellulose, gum arabic, polyvinyl alcohols, turpentine, bran oil, bentonite and ligninsulfonates.

To improve the flowability of a solid
25 product, there may be used adjuvants such as waxes, stearates and alkyl phosphates.

Adjuvants such as naphthalenesulfonic acid condensation products and polycondensates of phosphates

may be used as a peptizer for dispersible products.

Adjuvants such as silicone oil may also be used as a defoaming agent.

The content of the active ingredient may be varied according to the need, in a range of 0.01 to 80 parts by weight per 100 parts by weight of the preparation. In dusts or granules, the suitable content thereof is from 0.01 to 50% by weight. In emulsifiable concentrate and flowable wettable powder, too, the suitable content is from 0.01 to 50% by weight.

The agrohorticultural insecticide of the present invention is used to control a variety of insect pests in the following manner. That is, it is applied to a crop on which the insect pests are expected to appear or a site where appearance of the insect pests is undesirable, as it is or after being properly diluted with or suspended in water or the like, in an amount effective for control of the insect pests.

The applying dosage of the agrihorticultural insecticide of the present invention is varied depending upon various factors such as a purpose, insect pests to be controlled, a growth state of a plant, tendency of insect pests appearance, weather, environmental conditions, a preparation form, an application method, an application site and an application time. It may be properly chosen in a range of 0.1 g to 10 kg (in terms of active ingredient compound) per

10 ares depending upon purposes.

The agrihorticultural insecticide of the present invention may be used in admixture with other agricultural and horticultural disease or pest control-
 5 lers in order to expand both spectrum of controllable diseases and insect pest species and the period of time when effective applications are possible or to reduce the dosage.

Next, typical formulation examples and test
 10 examples of the invention are presented below. The present invention is by no means limited by these examples.

In the formulation examples, the term "parts" means "parts by weight".

15 Formulation Example 1

Each compound listed in Tables 2 to 12	50 parts
Xylene	40 parts
Mixture of polyoxyethylene nonylphenyl ether and calcium alkylbenzenesulfonate	10 parts

20 An emulsifiable concentrate was prepared by mixing uniformly the above ingredients to effect dissolution.

Formulation Example 2

Each compound listed in Tables 2 to 12	3 parts
25 Clay powder	82 parts
Diatomaceous earth powder	15 parts

A dust was prepared by mixing uniformly and grinding the above ingredients.

Formulation Example 3

	Each compound listed in Tables 2 to 12	5 parts
5	Mixed powder of bentonite and clay	90 parts
	Calcium ligninsulfonate	5 parts

Granules were prepared by mixing the above ingredients uniformly, and kneading the resulting mixture together with a suitable amount of water,
10 followed by granulation and drying.

Formulation Example 4

	Each compound listed in Tables 2 to 12	20 parts
	Mixture of kaolin and synthetic	
	high-dispersion silicic acid	75 parts
15	Mixture of polyoxyethylene nonylphenyl	
	ether and calcium alkylbenzenesulfonate	5 parts

A wettable powder was prepared by mixing uniformly and grinding the above ingredients.

Test Example 1: Insecticidal effect on diamond back
20 moth (Plutella xylostella)

Adult diamond back moths were released and allowed to oviposit on a Chinese cabbage seedling. Two days after the release, the seedling having the eggs deposited thereon was immersed for about 30 seconds in
25 a liquid chemical prepared by diluting a preparation

immersion, the hatched insects were counted. The mortality was calculated according to the following equation and the insecticidal effect was judged according to the criterion shown below. The test was carried out with triplicate groups of 10 insects.

10

A --- Mortality 100%

B --- Mortality 99-90%

C --- Mortality 89-80%

D --- Mortality 79-50%

- 15

Q1-12, Q1-41, Q1-42, Q4-6, Q4-8, Q4-12, Q4-45, Q6-6,
20 Q6-8, Q8-1, Q8-2, Q8-3, Q8-5, Q8-11, Q8-13, Q8-14, Q8-
15, Q8-53, Q9-15, Q10-7, Q15-1, Q26-1, Q26-11, Q26-29,
Q26-30, Q26-31, Q26-32, Q26-33, Q26-43, Q27-29, Q27-30,
Q27-31, Q27-32, Q27-81, Q27-98, Q27-124, Q27-125, Q27-

126, Q27-127, Q27-128, Q27-129, Q27-130, Q27-131, Q27-132, Q27-133, Q27-134, Q27-135, Q27-136, Q27-137, Q27-138, Q27-139, Q27-140, Q27-141, Q27-142, Q27-143, Q27-144, Q27-145, Q27-150, Q27-151, Q27-153, Q27-155, Q27-164, Q27-230, Q27-231, Q27-232, Q27-233, Q27-234, Q27-235, Q27-236, Q27-238, Q27-239, Q27-240, Q27-241, Q27-242, Q27-243, Q27-244, Q27-245, Q27-246, Q27-247, Q27-248 to Q27-265, Q27-267, Q27-268, Q27-270, Q27-273 to Q27-280, Q27-282 to Q27-284, Q27-289, Q27-297, Q27-298, Q27-305 to Q27-309, Q27-316 to Q27-318, Q27-323 to Q27-327, Q27-332, Q27-334, Q27-335, Q27-336 to Q27-339, Q27-347, Q27-352, Q27-353, Q27-354, Q27-355, Q27-359, Q27-360, Q27-378, Q27-384, Q27-387, Q27-388, Q27-391, Q27-393, Q27-396, Q27-397, Q27-402 to Q27-405, Q27-407, Q27-413, Q27-414, Q27-439, Q27-449 to Q27-457, Q27-459 to Q27-469, Q32-11, Q32-19, Q33-1, Q33-10, Q34-30, Q34-46 and Q42-1.

Test Example 2: Insecticidal effect on Common cutworm
(Spodoptera litura)

20 A piece of cabbage leaf (cultivar; Shikidori) was immersed for about 30 seconds in a liquid chemical prepared by diluting a preparation containing each compound listed in Tables 2 to 12 as an active ingredient to adjust the concentration to 500 ppm.

25 After air-dryness, it was placed in a plastic Petri dish with a diameter of 9 cm and inoculated with second-instar larvae of common cutworm, after which the

dish was closed and then allowed to stand in a room thermostatted at 25°C. Eight days after the inoculation, the dead and alive were counted. The mortality was calculated according to the following equation and the insecticidal effect was judged according to the criterion shown in Test Example 1. The test was carried out with triplicate groups of 10 insects.

$$\text{Corrected mortality(\%)} = \frac{\text{Number of alive larvae in untreated group} - \text{Number of alive larvae in treated group}}{\text{Number of alive larvae in untreated group}} \times 100$$

- 10 In the test mentioned above, the compounds which exhibited an activity ranking B or higher were as follows:
- Q26-1, Q26-11, Q26-29, Q26-30, Q26-31, Q26-32, Q26-33, Q26-43, Q27-29, Q27-30, Q27-31, Q27-32, Q27-81, Q27-98,
- 15 Q27-124, Q27-125, Q27-126, Q27-127, Q27-128, Q27-129, Q27-130, Q27-131, Q27-132, Q27-133, Q27-134, Q27-135, Q27-136, Q27-137, Q27-138, Q27-139, Q27-140, Q27-141, Q27-142, Q27-143, Q27-144, Q27-145, Q27-150, Q27-151, Q27-152, Q27-153, Q27-155, Q27-164, Q27-230, Q27-231,
- 20 Q27-232, Q27-233, Q27-234, Q27-235, Q27-236, Q27-238, Q27-239, Q27-240, Q27-241, Q27-242, Q27-243, Q27-244, Q27-245, Q27-246, Q27-247, Q27-248 to Q27-265, Q27-267, Q27-268, Q27-270, Q27-273 to Q27-280, Q27-282 to Q27-284, Q27-289, Q27-297, Q27-298, Q27-305 to Q27-309,

Q27-316 to Q27-318, Q27-323 to Q27-327, Q27-332, Q27-334, Q27-335, Q27-336 to Q27-339, Q27-347, Q27-352, Q27-353, Q27-354, Q27-355, Q27-359, Q27-360, Q27-378, Q27-384, Q27-387, Q27-388, Q27-397, Q27-402 to Q27-405,
5 Q27-407, Q27-413, Q27-414, Q27-439, Q27-459, Q27-466, Q32-19 and Q34-46.

Test Example 3: Insecticidal effect on rice leafroller
(Cnaphalocrosis medinalis)

The lamina of a rice plant at the 6 to 8 leaf
10 stage was immersed for about 30 seconds in a liquid chemical prepared by diluting a preparation containing each compound listed in Tables 2 to 12 as an active ingredient to adjust the concentration to 500 ppm. After air-dryness, the lamina was placed in a plastic
15 Petri dish with a diameter of 9 cm whose bottom had been covered with a wetted filter paper. The lamina was inoculated with third-instar larvae of rice leafroller, after which the dish was allowed to stand in a room thermostatted at 25°C and having a humidity
20 of 70%. Four days after the inoculation, the dead and alive were counted and the insecticidal effect was judged according to the criterion shown in Test Example 1. The test was carried out with triplicate groups of 10 insects.

25 In the test mentioned above, compounds which exhibited an activity ranking B or higher were as follows:

Q26-1, Q26-29, Q26-30, Q26-31, Q26-32, Q26-33, Q27-29,
Q27-30, Q27-31, Q27-32, Q27-81, Q27-98, Q27-124, Q27-
125, Q27-126, Q27-127, Q27-128, Q27-129, Q27-130, Q27-
131, Q27-132, Q27-133, Q27-134, Q27-135, Q27-136, Q27-
5 137, Q27-138, Q27-139, Q27-140, Q27-141, Q27-142, Q27-
143, Q27-144, Q27-145 and Q27-164.

2025-10-10 10:10:10